

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/105675/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Blokland, Gabriëlla A.M., del Re, Elisabetta C., Mesholam-Gately, Raquelle I., Jovicich, Jorge, Trampush, Joey W., Keshavan, Matcheri S., DeLisi, Lynn E., Walters, James ORCID: <https://orcid.org/0000-0002-6980-4053>, Turner, Jessica A., Malhotra, Anil K., Lencz, Todd, Shenton, Martha E., Voineskos, Aristotle N., Rujescu, Dan, Giegling, Ina, Kahn, René S., Roffman, Joshua L., Holt, Daphne J., Ehrlich, Stefan, Kikinis, Zora, Dazzan, Paola, Murray, Robin M., Di Forti, Marta, Lee, Jimmy, Sim, Kang, Lam, Max, Wolthusen, Rick P.F., de Zwarte, Sonja M.C., Walton, Esther, Cosgrove, Donna, Kelly, Sinead, Maleki, Nasim, Osiecki, Lisa, Picchioni, Marco M., Bramon, Elvira, Russo, Manuela, David, Anthony S., Mondelli, Valeria, Reinders, Antje A.T.S., Falcone, M. Aurora, Hartmann, Annette M., Konte, Bettina, Morris, Derek W., Gill, Michael, Corvin, Aiden P., Cahn, Wiepke, Ho, New Fei, Liu, Jian Jun, Keefe, Richard S.E., Gollub, Randy L., Manoach, Dara S., Calhoun, Vince D., Schulz, S. Charles, Sponheim, Scott R., Goff, Donald C., Buka, Stephen L., Cherkerzian, Sara, Thermenos, Heidi W., Kubicki, Marek, Nestor, Paul G., Dickie, Erin W., Vassos, Evangelos, Ciufolini, Simone, Reis Marques, Tiago, Crossley, Nicolas A., Purcell, Shaun M., Smoller, Jordan W., van Haren, Neeltje E.M., Touloupoulou, Timothea, Donohoe, Gary, Goldstein, Jill M., Seidman, Larry J., McCarley, Robert W. and Petryshen, Tracey L. 2018. The genetics of endophenotypes of neurofunction to understand schizophrenia (GENUS) consortium: a collaborative cognitive and neuroimaging genetics project. *Schizophrenia Research* 195 , pp. 306-317. 10.1016/j.schres.2017.09.024 file

Publishers page: <http://dx.doi.org/10.1016/j.schres.2017.09.024>
<<http://dx.doi.org/10.1016/j.schres.2017.09.024>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<https://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright

information services
gwasanaethau gwybodaeth



holders.

Supplementary Material for:
The Genetics of Endophenotypes of Neurofunction to Understand Schizophrenia (GENUS)
Consortium: A collaborative cognitive and neuroimaging genetics project

| | |
|---|-----------|
| Sample Descriptions..... | 2 |
| CAMH — Center for Addiction and Mental Health | 2 |
| CATIE — Clinical Antipsychotic Trials of Intervention Effectiveness | 3 |
| CIDAR/VA — Boston CIDAR Study / Veterans Affairs Healthcare System | 4 |
| COGS-UK — Cognition in Schizophrenia, Cardiff University, UK | 6 |
| GAP — Genetics and Psychosis study | 7 |
| IMH-SIGNRP — Institute of Mental Health, Singapore Imaging Genetics and Neuropsychological Research in Psychosis | 8 |
| IMH-STCRP — Institute of Mental Health, Singapore Translational and Clinical Research in Psychosis.... | 9 |
| KCL-MFS — King’s College London - Maudsley Family Study | 10 |
| KCL-MTS — King’s College London - Maudsley Twin Study | 12 |
| L&R — Language and Risk in Schizophrenia | 13 |
| MCIC — Mind Clinical Imaging Consortium | 15 |
| MGH — Massachusetts General Hospital | 17 |
| NEFS — New England Family Study..... | 18 |
| PAGES — Phenomics and Genomics Sample | 22 |
| PHRS — Pittsburgh High Risk Study..... | 23 |
| TCD/NUIG — Trinity College Dublin and National University of Ireland, Galway | 24 |
| UMCU-SZ1 — University Medical Centre Utrecht, Schizophrenia study 1 | 25 |
| UMCU-SZ2 — University Medical Centre Utrecht, Schizophrenia study 2 | 26 |
| ZHH — Zucker Hillside Hospital | 27 |
| References | 29 |

Sample Descriptions

CAMH — Center for Addiction and Mental Health

(Nazeri et al., 2013; Wheeler et al., 2014)

Recruitment. Patients and controls were recruited at the Center for Addiction and Mental Health (CAMH) in Toronto, Canada, via referrals, study registries, and advertisements. All clinical assessments occurred at CAMH and MRI scans were performed at a nearby general hospital in Toronto. Community-dwelling outpatients with schizophrenia were individually matched based on age, sex, handedness, and highest parental years of education to healthy controls. Controls form part of the Cognitive Genomics Consortium wave 2 (COGENT2) dataset (Trampush et al., 2017).

Inclusion/Exclusion Criteria. All participants were between 18 and 85 years of age. All participants received urine toxicology screens and anyone with current substance abuse or any history of substance dependence was excluded. Individuals with previous head trauma with loss of consciousness or neurological disorders were also excluded. For controls, a history of a primary psychotic disorder in first-degree relatives was also an exclusion criterion.

Clinical/Neuropsychological Assessment. All participants were interviewed by a psychiatrist and were administered the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I; First et al., 1995b) to determine diagnosis and duration of illness. The Positive and Negative Syndrome Scale (PANSS; Kay et al., 1987) and Clinical Global Impression Scale (CGI-S; Guy, 2000) were administered to further characterize illness symptoms. Burden of comorbid physical illness was measured by the Cumulative Illness Rating Scale for Geriatrics (Miller et al., 1992). Medication histories were initially recorded based on self-reports, and then verified either by the patient's treating psychiatrist or chart review. All participants underwent a battery of cognitive tests administered over approximately 1.5 hours. Premorbid IQ was measured using the Wechsler Test for Adult Reading (WTAR; Wechsler, 2001). The cognitive battery consisted of the subtests Digit Symbol Coding, Semantic Verbal Fluency, Digit Span, List Learning, Story Memory, and Figure Recall from the Repeated Battery for the Assessment of Neuropsychological Status (RBANS; Randolph et al., 1998), as well as the Trail Making Test-Part A and B (Reitan, 1958) and Controlled Oral Word Association Test (COWAT) of Phonemic Verbal Fluency (letters: F, A, S) (Reitan, 1985) to measure processing speed, University of Maryland Letter-Number Span (Gold et al., 1997) to measure verbal working memory, Stroop test to measure executive function (Stroop, 1935), Grooved Pegboard (Lafayette Instrument Company -- Trites, 1989) and Finger Tapping (Ruff and Parker, 1993) to assess psychomotor speed.

MRI Acquisition. T1-weighted structural scans (Echo Time [TE] = 5.3 ms; Inversion Time [TI] = 300 ms; Repetition Time [TR] = 12.3 ms; Flip angle = 20°; Field of View [FOV] = 20 cm; Spatial resolution = 0.78 x 0.78 x 1.5 mm; matrix = 256 x 256 mm; Slice acquisition direction = axial, AC-PC oblique) and Diffusion Tensor Imaging (TE = 85.5 ms; TR = 15000 ms; Flip angle = 90°; FOV = 33 cm; Spatial resolution = 2.6 x 2.6 x 2.6 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial, oblique; number of diffusion-encoding gradient directions = 23; number of b0 volumes = 2; b-value = 1000 s/mm²) were acquired on a 1.5 Tesla General Electric (GE) Echosped scanner using an 8-channel head coil.

Genotyping. DNA obtained from whole blood was genotyped on the Illumina OmniExpress array.

Ethics/Consent. The study was approved by the Research Ethics Board of CAMH and all participants provided informed written consent.

Acknowledgements. Data acquisition was supported by the Canadian Institutes of Health Research, Ontario Mental Health Foundation, the Brain and Behavior Research Foundation (formerly NARSAD), the Centre for Addiction and Mental Health (CAMH), and the CAMH Foundation through the Kimel Family, Koerner New Scientist Award, and Paul E Garfinkel New

Investigator Catalyst Award. The investigators acknowledge the assistance of David J. Rotenberg and Dr. Daniel Felsky.

CATIE — Clinical Antipsychotic Trials of Intervention Effectiveness

(Keefe et al., 2006; Lieberman, 2007; McClay et al., 2011; Stroup et al., 2003; Sullivan et al., 2008)

Recruitment. Patients were recruited from 57 clinical settings across the United States for a multiphase randomized controlled trial of antipsychotic medications, including the second-generation drugs, olanzapine, quetiapine, risperidone, and ziprasidone, compared with a midpotency first-generation drug, perphenazine. Patients were followed for up to 18 months. Patients who gave informed consent to participate in genetic studies are included in GENUS. Patients formed part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. Eligible patients were 18 to 65 years of age; had received a diagnosis of schizophrenia (not schizoaffective disorder), as determined on the basis of the SCID; and were able to take oral antipsychotic medication, as determined by the study doctor. Patients were excluded if they had received a diagnosis of schizoaffective disorder, mental retardation, or other cognitive disorders; had a history of serious adverse reactions to the proposed treatments; had had only one schizophrenic episode; had a history of treatment resistance, defined by the persistence of severe symptoms despite adequate trials of one of the proposed treatments or prior treatment with clozapine; were pregnant or breastfeeding; or had a serious and unstable medical condition. Patients with medical or psychiatric comorbidities those who require concomitant other medications are included. The patients who enroll in this study have chronic or recurrent schizophrenia. First episode patients and wholly treatment-refractory patients are excluded. First episode patients are excluded because of their high rates of response to antipsychotic medications at relatively low doses. Treatment-refractory patients are excluded because their severe illness may preclude detection of differential effectiveness that would be apparent in treatment-responsive patients.

Clinical/Neuropsychological Assessment. Schizophrenia patients were diagnosed using the SCID (First et al., 1994). Clinical rating scales include the PANSS (Kay et al., 1987), Calgary Depression Scale for Schizophrenia (CDSS; Addington et al., 1992), and Clinical Global Impression Scale (CGI-S; Guy, 2000). Premorbid IQ was measured using the Reading subtest from the Wide Range Achievement Test, Third Edition (WRAT-3; Wilkinson, 1993). Processing speed was measured using the Wechsler Adult Intelligence Scale, Revised Version (WAIS-R) Digit Symbol subtest (Wechsler, 1981), Semantic Verbal Fluency (categories: animals, fruits, vegetables) (Benton and Hamsher, 1978), and COWAT Phonemic Verbal Fluency (letters: F, A, S) (Reitan, 1985); attention/vigilance was measured using the Continuous Performance Test-Identical Pairs version (CPT-IP; Cornblatt et al., 1988); spatial and verbal working memory were assessed using the computerized Spatial Delayed Response Task (Lyons-Warren et al., 2004) and University of Maryland Letter-Number Span (Gold et al., 1997), respectively. Verbal learning and memory was assessed with the Hopkins Verbal Learning Test (HVLT; Brandt, 1991). Executive function was assessed using the Wisconsin Card Sorting Test (WCST; Heaton et al., 1993) and the Mazes subtest from the Wechsler Intelligence Scale for Children, Third Edition (WISC-III; Wechsler, 1991). Grooved Pegboard (Lafayette Instrument Company -- Trites, 1989) was used to measure psychomotor speed and the Facial Emotion Discrimination Test (Kerr and Neale, 1993) measured social cognition.

MRI Acquisition. Not applicable.

Genotyping. DNA obtained from whole blood was genotyped on the Affymetrix 500K array and Perlegen's custom 164K chip. Genotype data are available from the NIMH Repository and Genomics Resource (<http://www.nimhgenetics.org>).

Ethics/Consent. All participants or their legal guardians gave written informed consent and the institutional review board at each site approved the study.

Acknowledgements. The principal investigators were Jeffrey A. Lieberman, M.D., T. Scott Stroup, M.D., M.P.H., and Joseph P. McEvoy, M.D. The CATIE trial was funded by a grant from the National Institute of Mental Health (N01 MH900001) along with MH074027 (PI PF Sullivan). Genotyping was funded by Eli Lilly and Company.

Conflicts of interest. Dr. Sullivan reports receiving research funding from Eli Lilly in connection with the CATIE project. Dr. Stroup reports having received research funding from Eli Lilly and consulting fees from Janssen Pharmaceutica, GlaxoSmithKline, and Bristol-Myers Squibb. Dr. Lieberman reports having received research funding from AstraZeneca Pharmaceuticals, Bristol-Myers Squibb, GlaxoSmithKline, Janssen Pharmaceutica, and Pfizer and consulting and educational fees from AstraZeneca Pharmaceuticals, Bristol-Myers Squibb, Eli Lilly, Forest Pharmaceuticals, GlaxoSmithKline, Janssen Pharmaceutica, Novartis, Pfizer, and Solvay.

CIDAR/VA — Boston CIDAR Study / Veterans Affairs Healthcare System

(Clemm von Hohenberg et al., 2014; del Re et al., 2014; Kikinis et al., 2010; Kikinis et al., 2015; Pasternak et al., 2012; Quan et al., 2013; Seitz et al., 2016)

Recruitment. Participants were recruited through the Boston CIDAR study (www.bostoncidar.org) and the Veterans Affairs Healthcare System. The participants included in the GENUS dataset consisted of clinical high-risk individuals with prodromal symptoms who also have a family history of schizophrenia (included in the FHR group), individuals within the first episode of schizophrenia, and individuals with chronic schizophrenia, as well as group-matched healthy control participants. Patients and FHR participants were recruited by referrals from clinicians or through local hospitals and clinics, and controls were recruited through newspaper and website advertisements. Patient and control participants of European ancestry formed part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. Exclusion criteria for controls included any current or past major DSM-IV-TR Axis I disorder, developmental disorders, psychiatric hospitalizations, prodromal symptoms, schizotypal or other Cluster A personality disorders, a first degree relative with psychosis, or current or past use of antipsychotics. Sleeping aids or anxiolytic agents for occasional use were acceptable, as well as other past psychotropic medication use if not within the prior 6 months. Exclusion criteria for all participants included sensory-motor handicaps, neurological disorders, medical illnesses that significantly impair neurocognitive function, diagnosis of intellectual disability (IQ<70), not fluent in English, DSM-IV substance abuse within the past month or dependence within the past 3 months, current suicidality, or history of electroconvulsive therapy for controls or electroconvulsive therapy within the past 5 years for patients. All first episode patients met DSM-IV-TR criteria for schizophrenia, schizoaffective disorder or schizophreniform disorder, and none had more than one year of continuous antipsychotic treatment. Chronic patients met DSM-IV-TR criteria for schizophrenia or schizoaffective disorder and had illness duration of at least five years. Inclusion criteria for clinical high-risk participants was predominantly presence of ‘late prodromal syndromes’ as classified by the Criteria of Prodromal States (Woods et al., 2001) based on the Scale of Prodromal Symptoms (Miller et al., 2003; Miller et al., 1999). Of the clinical high-risk participants, 14 participants fulfilled the GENUS definition of FHR. Controls were drawn from the same geographic bases with comparable age, gender, race and ethnicity, handedness, and parental socioeconomic status evaluated using the Hollingshead two-factor index (Hollingshead, 1975).

Clinical/Neuropsychological Assessment. Trained and skilled interviewers and neuropsychological testers conducted all clinical and cognitive assessments. DSM-IV diagnoses were based on interviews with the Structured Clinical Interview for DSM-IV-TR (SCID), Research Version (First et al., 2002b) and information from patient medical records. The Scale for the Assessment of Negative Symptoms (SANS; Andreasen, 1983) and the Scale for the Assessment

of Positive Symptoms (SAPS; Andreasen, 1984) were used to measure symptom severity. The Global Assessment of Functioning (GAF; American Psychiatric Association, 2000) and CGI-S (Guy, 2000) were used to measure overall functioning. Participants were administered a large neurocognitive test battery including tests estimating premorbid IQ (the Reading subtest from the Wide Range Achievement Test-4 (WRAT-4; Wilkinson and Robertson, 2006a) and current IQ (the Vocabulary and Block Design subtests of the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler, 1999). The Brief Assessment of Cognition in Schizophrenia (BACS) Symbol Coding test (Keefe et al., 2008), the Trail Making Test-Part A (Reitan, 1958), and Semantic Verbal Fluency (MCCB Category Fluency; category: animals) (Benton and Hamsher, 1978; Kern et al., 2008; Nuechterlein et al., 2008) were used for processing speed; the Continuous Performance Test-Identical Pairs version (Cornblatt et al., 1988) was used to measure attention/vigilance; the Wechsler Memory Scale, Third Edition (WMS-III) Spatial Span subtest (Wechsler, 1997b) and University of Maryland Letter-Number Span (Gold et al., 1997) were used for assessing working memory. The ‘vigilance’, ‘working memory’ and ‘working memory plus interference’ measures from the Seidman Auditory Continuous Performance Test battery (Auditory CPT; (Seidman et al., 1998; Seidman et al., 2006). The Hopkins Verbal Learning Test-Revised (HVLT-R; Brandt and Benedict, 1997) and the WMS-III Logical Memory subtest (Wechsler, 1997b) assessed verbal learning and memory. The Brief Visuospatial Memory Test-Revised (BVM-T-R; Benedict, 1997) assessed visual learning. Executive function was measured using the Mazes subtest of the Neuropsychological Assessment Battery (NAB; Stern and White, 2003) and the computerized WCST (Heaton, 2003; Heaton et al., 1993; Nestor et al., 1993). Finally, the Proverbs subtest from the Delis-Kaplan Executive Function System (D-KEFS; (Delis et al., 2001; Delis et al., 2004)) was administered to assess verbal ability, and Grooved Pegboard (Lafayette Instrument Company -- Trites, 1989) and Finger Tapping (Ruff and Parker, 1993) were administered to assess psychomotor speed. Social cognition was assessed using the Mayer–Salovey–Caruso Emotional Intelligence Test (MSCEIT; Mayer et al., 2003).

MRI Acquisition. T1-weighted structural scans of schizophrenia and age-gender-handedness matched controls were acquired at Brigham and Women’s Hospital, Harvard University (TE = 3.0 ms; TI = 600 ms; TR = 7.8 ms; Flip angle = 10°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.0 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 78 ms; TR = 17000 ms; Flip angle = 90°; FOV = 24 cm; Spatial resolution = 1.67 x 1.67 x 1.7 mm; matrix = 144 x 144 mm; Slice acquisition direction = axial, AC-PC; number of diffusion-encoding gradient directions = 51; number of b0 volumes = 8; b-value = 900 s/mm²) were acquired on a 3 Tesla GE Signa HDxt Echospeed scanner using an 8-channel head coil.

T1-weighted structural scans of FHR participants and age-gender-handedness matched controls were acquired at Massachusetts General Hospital, Martinos Center, Harvard University, on a 3 Tesla Siemens Trio Tim scanner using a 32-channel head coil (TE = 3.48 ms; TI = 900 ms; TR = 2530 ms; Flip angle = 9°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.33 mm; matrix = 256 x 256 x 128 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 84 ms; TR = 9400 ms; Flip angle = 90°; FOV = 25.6 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 60; number of b0 volumes = 10; b-value = 700 s/mm²).

Genotyping. DNA was obtained from whole blood or saliva and genotyped on the Illumina OmniExpress array.

Ethics/Consent. The study was approved by the local IRB committees at Harvard Medical School, Beth Israel Deaconess Medical Center, Massachusetts General Hospital, Brigham and Women’s Hospital and the Veteran Affairs Boston Healthcare System (Brockton campus). All study participants, or legal guardians for those under 18 years of age, gave written informed consent and received payment for participation.

Acknowledgements. Data acquisition was supported by the National Institute of Mental Health (NIMH) of the National Institutes of Health (NIH) under Award Numbers P50 MH080272 (RW

McCarley), U01MH081928 (LJ Seidman), R21MH106793 (Z Kikinis), and a NARSAD Young Investigator Award (Z Kikinis). Genotyping was funded by a Massachusetts General Hospital Executive Committee on Research Interim Support Fund award (TL Petryshen). The investigators thank all subjects for their participation in the study, the clinical and data management staff from the Boston CIDAR study and the Commonwealth Research Center (Dr. Matcheri Keshavan, Dr. Joanne Wojcik, Ann Cousins, Dr. Michelle Friedman-Yakoobian, Dr. Anthony J Giuliano, Andrea Gnong Granato, Lauren Gibson, Sarah Hornbach, Julia Schutt, Dr. Kristy Klein, Dr. Maria Hiraldo, Dr. Grace Francis, Corin Pilo, Rachael Serur, Grace Min, Alison Thomas, and Molly Franz), Tamara Tasoff, Dr. Beril Yaffe, and Danbee Kim for electrophysiology assistance (Harvard), and Kimberly Chambert (Broad Institute of MIT and Harvard), Patience Gallagher, Dr. Stephen Haddad, Brian Galloway, and Jenna Tarasoff (Massachusetts General Hospital) for genotyping assistance.

COGS-UK — Cognition in Schizophrenia, Cardiff University, UK

Recruitment. Patients were ascertained from psychiatric outpatient units, via voluntary mental health organizations and by public advert and entered the study following a telephone screening interview and confirmation of a clinical diagnosis of a non-affective psychosis by the treating clinical team. The sample consisted of white Caucasians, born in the UK and with British ancestry.

Inclusion/Exclusion Criteria. Patients were screened to exclude substance-induced psychotic disorder or psychosis due to a general medical condition. All patients were diagnosed with schizophrenia, schizophreniform or schizoaffective diagnoses.

Clinical/Neuropsychological Assessment. A semi-structured interview was administered to all patients by trained psychiatrists or psychologists. The comprehensive clinical phenotype battery included the Schedules for Clinical Assessment in Neuropsychiatry (SCAN; Wing et al., 1990)) sections on psychosis and affective symptoms, substance use, and family history as well as demographic details and a separate cognitive battery. Diagnosis was made according to DSM-IV criteria by experienced psychiatrists accessing the clinical interview data and case notes when appropriate. Clinical rating scales administered included SANS/SAPS (Andreasen, 1983, 1984), CDSS (Addington et al., 1992), and GAF (American Psychiatric Association, 1994). Premorbid IQ was estimated using the National Adult Reading Test (NART; (Nelson, 1982; Nelson and Willison, 1991)). Cognitive testing was performed using the MATRICS Consensus Cognitive Battery (MCCB; (Kern et al., 2008; Nuechterlein et al., 2008)). The MCCB evaluates seven domains of cognitive function including: speed of processing using the BACS Symbol Coding subtest (Keefe et al., 2008), Trail Making Test—Part A (Reitan, 1958), and Semantic Verbal Fluency (MCCB Category Fluency; category: animals) (Benton and Hamsher, 1978; Kern et al., 2008; Nuechterlein et al., 2008); attention using the Continuous Performance Test—Identical Pairs (CPT-IP; Cornblatt et al., 1988); working memory using the WMS-III Spatial Span subtest (Wechsler, 1997b) and University of Maryland Letter-Number Span (Gold et al., 1997) (spatial and letter-number span); verbal learning using the HVLT-R (Brandt and Benedict, 1997); visual learning using the BVMT-R (Benedict, 1997); reasoning and problem solving using the NAB Mazes subtest (Stern and White, 2003); and social cognition using the MSCEIT (Mayer et al., 2003). Finally, the D-KEFS Proverbs subtest (Delis et al., 2001; Delis et al., 2004) was administered to assess verbal ability.

MRI Acquisition. Not applicable.

Genotyping. DNA obtained from whole blood from a proportion of patients was genotyped on the Illumina Infinium OmniExpressExome-8 array.

Ethics/Consent. The UK Multi-center Research Ethics Committee (MREC) approved the study, as did relevant local research and development committees. All participants provided valid informed consent for the study.

Acknowledgements. The investigators wish to acknowledge Dr. Antonio F. Pardiñas and Amy Lynham for their assistance.

GAP — Genetics and Psychosis study

(O'Connor et al., 2013; Reis Marques et al., 2014; Theleritis et al., 2014; Vassos et al., 2017)

Recruitment. First episode psychosis patients were recruited as part of the National Institute of Health Research (NIHR) Biomedical Research Centre (BRC) Genetics and Psychosis study. Patients were approached if they appeared to meet DSM-IV criteria for schizophrenia or related disorder, or affective disorder with psychotic features and presented to one of 3 south London boroughs' geographical catchment area-based adult mental health services services of the South London and Maudsley NHS Foundation Trust (SLaM). During the same period a healthy control group was recruited from the local population living in the area and served by the SLaM Trust by means of internet and newspaper advertisements, and distribution of leaflets at train stations, shops and job centers. Particular attention was directed at attempting to obtain a control sample similar to the patient sample in age, gender, ethnicity, educational qualifications and employment status (Di Forti et al., 2009).

Inclusion/Exclusion Criteria. Between August 2008 and July 2011, patients aged between 18–65 years presenting for the first time to psychiatric services with a functional psychotic illness (ICD-10: F20-29, F30-34 excluding coding F1x.0 for acute intoxication; World Health Organization, 1992)). Inclusion criteria required that patients have 7 or more consecutive days of psychotic symptom(s) and were presenting to services for the first time with these symptoms. Patients were approached as soon as possible (up to 3 months after first contact). Exclusion criteria were: presence of an organic psychosis (i.e. known organic cause for psychosis), a moderate or severe developmental learning disability (as defined by ICD-10 F70-73), pregnancy, history of a medical or physiological cause of gonadal dysfunction (including hypothyroidism or other endocrine or metabolic disorder), vascular disorders, neurological disorders, antipsychotic treatment longer than 30 days (to minimize the effect of antipsychotics on BDNF levels), poor English fluency, and history of contact with health services for psychosis beyond the previous 6 months. Participants completed all measures within 6 months of first presentation with services. At 12 months after initial assessments, some research measures were repeated via clinical notes or face-to-face interview when available. Information about employment, relationships, living status and medication, was collected through interview with the patient and searching clinical records. Diagnosis was made according to DSM-IV criteria using the Operational Criteria (OPCRIT; McGuffin et al., 1991) based on the clinical notes for the month after first contact with psychiatric services for psychosis. Raters were experienced researchers who were extensively trained and demonstrated adequate inter-rater reliability on clinical and neuropsychological assessment. All diagnoses were carried out by qualified psychiatrists and clinical researchers, subject to comprehensive training and inter-rater reliability testing.

Clinical/Neuropsychological Assessment. Socio-demographic data (age, gender, self-reported ethnicity, level of education attainment) were collected using the Medical Research Council Social Scale (Mallett et al., 2002). Medication histories were completed using information from patients and electronic clinical records. The GAF (American Psychiatric Association, 1994) was used to rate both severity of symptoms and disability. Two separate ratings on the GAF were assigned based on dimensions of psychiatric symptoms and psychological, social and occupational function. To rate specific psychotic symptoms, scored on a clinical interview, the PANSS was used (Kay et al., 1987). Neuropsychological assessments included the following measures: premorbid IQ from the WTAR (Wechsler, 2001) or NART (Nelson, 1982; Nelson and Willison, 1991; Russell et al., 2000)); spatial and verbal working memory using the WMS-III Spatial Span subtest (Wechsler, 1997b) and the Digit Span subtest from the Wechsler Adult Intelligence Scale, Third Edition (WAIS-III; Wechsler, 1997a)) and the Spatial Working Memory subtest from the Cambridge Neuropsychological Test Automated Battery (CANTAB; Cambridge Cognition, 2003a)); attention/processing speed using Trail Making Test—Part A (Reitan, 1958), WAIS-III Digit Symbol (Wechsler, 1997a), Semantic Verbal Fluency (categories: animals, body

parts, fruits) and Phonemic Verbal Fluency (letters F, A, S); verbal learning and memory was assessed using WMS-III Logical Memory (Wechsler, 1997b); visual memory using WMS-III Visual Reproduction; executive function was assessed using Trail Making Test—Part B (Reitan, 1958) and CANTAB Tower of London (Cambridge Cognition, 2003a). Additionally, the WAIS-III subtests Block Design, Matrix Reasoning, and Information, along with Digit Symbol and Digit Span were administered to allow for a calculation of total current IQ (Wechsler, 1997a).

MRI Acquisition. T1-weighted structural scans (TE = 2.85 ms; TI = 650 ms; TR = 6.99 ms; Flip angle = 8°; FOV = 26 cm; Spatial resolution = 1.0 x 1.0 x 1.2 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 104.8 ms; TR = 15000 ms; Flip angle = 90°; FOV = 30.7 cm; Spatial resolution = 2.4 x 2.4 x 2.4 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 32; number of b0 volumes = 4; b-value = 1300 s/mm²) were acquired on a 3 Tesla GE Signa HDx scanner using an 8-channel head coil.

Genotyping. DNA from a proportion of participants was extracted from blood or cheek swabs (80% and 20% of the participants, respectively) and genotyped on the Illumina HumanCore-24 Exome BeadChip.

Ethics/Consent. Ethical approval was obtained from the local research ethics committee. All study participants gave full informed consent. After hearing a complete description of the study and having the opportunity to ask questions, participants gave written informed consent.

Acknowledgements. The GAP study was supported by the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. Dr. Dazzan's research is also supported by the Medical Research Foundation. Dr. Reis-Marques and Dr. Dazzan's research is supported by NARSAD. Dr. Reinders is supported by the Netherlands Organization for Scientific Research (NWO-VENI grant no. 451-07-009). Genotyping and basic QC was supported by Guy's and St. Thomas Charity Grant No.R080529 (E Vassos) and the Psychiatry Research Trust (R Murray). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health". The investigators are grateful for the support of all GAP researchers and Principal Investigators for their contribution and support of the GAP study, and gratefully acknowledge the help of the Genetic & Psychosis and Physical Health & Substance Use in First Episode Psychosis study teams and participants, South London & Maudsley Mental Health NHS Trust. The investigators wish to specifically acknowledge Dr. Carmine M. Pariante, Dr. Kie Woo Nam, and Heather Taylor.

Conflicts of interest. Dr. Robin Murray has received honoraria for lectures from Janssen, AstraZeneca, Lilly, Novartis and Bristol-Myers Squibb.

IMH-SIGNRP — Institute of Mental Health, Singapore Imaging Genetics and Neuropsychological Research in Psychosis

(Ho et al., 2016; Kuswanto et al., 2015)

Recruitment. Chinese ethnicity patients diagnosed with DSM-IV schizophrenia were recruited from the Institute of Mental Health, Singapore. Age- and gender-matched healthy controls were recruited from the staff population at the hospital, as well as from the community by advertisements.

Inclusion/Exclusion Criteria. Participants with history of any significant neurological illnesses such as seizure disorder, head trauma, or cerebrovascular accident were excluded. No subject met DSM-IV criteria for alcohol or other substance abuse in the preceding 3 months. Patients were maintained on a stable dose of antipsychotic medication for at least two weeks prior to the recruitment and did not have their medications withdrawn for the purpose of the study. Healthy controls were deemed not to suffer from any Axis I psychiatric disorder and had no history of any major neurological, medical illnesses, substance abuse or psychotropic medication use.

Clinical/Neuropsychological Assessment. All diagnoses were made by a psychiatrist using information obtained from the existing medical records, clinical history, mental status examination, interviews with the patients and their significant family members, as well as the administration of SCID-I, Patient Version (SCID-I/P; First et al., 1994). Healthy controls were screened using the SCID-I, Non-Patient Version (SCID-I/NP; First et al., 2002a). Psychotic symptom levels were assessed using the PANSS (Kay et al., 1987). Premorbid IQ was assessed using the Reading subscale of the WRAT-3 (Wilkinson, 1993). The Symbol Coding, Category Instances (category: supermarket items), COWAT Phonemic Verbal Fluency (Reitan, 1985) (letters: F and S), Digit Sequencing, List Learning, and Tower of London (Shallice, 1982) subtests from the BACS (Keefe et al., 2004) were administered to assess attention/processing speed, verbal working memory, verbal learning and memory, and executive function, respectively.

MRI Acquisition. T1-weighted structural scans (TE = 3.3 ms; TI = 860 ms; TR = 7.2 ms; Flip angle = 8°; FOV = 23 cm; Spatial resolution = 0.9 x 0.9 x 0.9 mm; matrix = 256 x 256 mm; Slice acquisition direction = axial) and Diffusion Tensor Imaging (TE = 56 ms; TR = 3725 ms; Flip angle = 90°; FOV = 23 cm; Spatial resolution = 0.9 x 0.9 x 3.9 mm; matrix = 112 x 109 mm; Slice acquisition direction = axial, AC-PC; number of diffusion-encoding gradient directions = 15; number of b0 volumes = 1; b-value = 800 s/mm²) were acquired on a 3 Tesla Philips Intera Achieva scanner using a SENSE 8-channel head coil.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on one of three arrays: Illumina Human OmniZhongHua-8, Illumina Human1M-Duo, or Affymetrix 6.0.

Ethics/Consent. This study was approved by the Institutional Review Board of the Institute of Mental Health, Singapore, as well as the National Neuroscience Institute, Singapore. Participants gave written informed consent to participate in the study after a detailed explanation of the study procedure.

Acknowledgements. Data acquisition was supported by the National Research Foundation Singapore under the National Medical Research Council Translational and Clinical Research Flagship Programme (grant no. NMRC/TCR/003/2008) for The Singapore Translational and Clinical Research in Psychosis, as well as research grants from the National Medical Research Council under the Centre Grant Programme (Institute of Mental Health, Singapore) (NMRC/CG/004/2013), the National Healthcare Group, Singapore (SIG/05004; SIG/05028), and the Singapore Bioimaging Consortium (RP C-009/2006). The investigators wish to acknowledge the contribution of Mingyuan Wang.

IMH-STCRP — Institute of Mental Health, Singapore Translational and Clinical Research in Psychosis

(Lam et al., 2014)

Recruitment. Patients and healthy controls were recruited as part of the Singapore Translational and Clinical Research in Psychosis. Patients were recruited from rehabilitation centers, community care centers across the country, out-patient clinics and in-patient wards, under purview of the Institute of Mental Health, Singapore. Data collection was completed in approximately 3 years. Controls were from the Singapore Prospective Study Program and were randomly sampled from the Singapore population. Participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. Inclusion criteria were: Chinese ethnicity, to ensure a genetically homogeneous sample; and completion of a minimum of 6 years of primary school education. Additional exclusion criteria precluded all participants with significant history of substance abuse, clinically significant neurological disease or injury, color blindness, and healthy participants with first-degree relatives suffering from schizophrenia or other psychotic disorders.

Clinical/Neuropsychological Assessment. Clinical and neuropsychological evaluations were carried out by psychometricians. Schizophrenia patients fulfilled Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) diagnostic criteria for schizophrenia based on the SCID-I (First et al., 2002b). Psychotic symptoms were assessed using the PANSS (Kay et al., 1987). Neuropsychological tests administered included the Matrix Reasoning subtest of the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler, 1999) as a measure of current IQ, Continuous Performance Tests-Identical Pairs (CPT-IP; Cornblatt et al., 1988) to measure attention/vigilance, 64-card WCST (WCST-64; Heaton et al., 1993) to measure executive function, Benton Judgment of Line Orientation Test (Benton et al., 1994) to measure visuospatial ability, and the BACS subtests Symbol Coding, Category Instances (categories: animals, fruits, vegetables), Digit Sequencing, List Learning, Tower of London, and Token Motor Task (Keefe et al., 2004; Keefe et al., 2008) to assess attention/processing speed, verbal working memory, verbal learning and memory, executive function, and psychomotor speed, respectively. Data including education, duration of illness, and medications were also collected.

MRI Acquisition. Not applicable.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on the Illumina HumanOmniZhongHua-8 BeadChip.

Ethics/Consent. All participants consented to participate in research procedures prior to data collection. Consent procedures adhered to the guidelines specified by the National Healthcare Group Domain Specific Review Board's (domain A, NHG DSRB) requirements for human subject research.

Acknowledgements. Data acquisition was supported by the National Research Foundation Singapore under the National Medical Research Council Translational and Clinical Research Flagship Programme (grant no. NMRC/TCR/003/2008) for The Singapore Translational and Clinical Research in Psychosis, as well as research grants from the National Medical Research Council under the Centre Grant Programme (Institute of Mental Health, Singapore) (NMRC/CG/004/2013), the National Healthcare Group, Singapore (SIG/05004; SIG/05028), and the Singapore Bioimaging Consortium (RP C-009/2006).

KCL-MFS — King's College London - Maudsley Family Study

(Psychosis Endophenotypes International Consortium et al., 2014; Touloupoulou et al., 2004; Touloupoulou et al., 2003)

Recruitment. Patients and their relatives were drawn from a larger sample of the Maudsley Family Study (Frangou et al., 1997a,b; Griffiths et al., 1998). The relative group comprised mainly first-degree relatives and some second degree-relatives. The latter were second degree to the proband but they themselves had a first-degree relative with schizophrenia who either did not want to participate in this aspect of the Maudsley Family Study or was deceased. All offspring were adults. Families were recruited by referrals of psychiatric clinics and various other organizations across the United Kingdom. All patients, the vast majority of whom were outpatients, met DSM-III-R criteria for schizophrenia or schizoaffective disorder and were on antipsychotic medication at the time of the assessment. Among the non-psychotic relatives were seven individuals who had had a single episode of major depression, and the following had a lifetime diagnosis: 10 recurrent depressive disorder, three schizotypal personality disorder, and one of each of bulimia nervosa, alcohol dependency and social phobia. All relatives were well at the time of the neuropsychological assessment. Control participants were recruited from a set of controls obtained for previous studies conducted at the Institute of Psychiatry, King's College, London, from members of staff at the Institute of Psychiatry and the Bethlem and Maudsley Hospitals and via advertisements in the local newspapers. Some participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. All participants spoke English as their primary language. The normal control sample consisted of healthy participants with no personal or family history, up to second degree, of psychotic illness. The following exclusion criteria were used for all participants: head trauma resulting in loss of consciousness, drug or alcohol abuse during the 12 months before testing, organic brain disease, English as non-primary language and IQ<70. An additional exclusion criterion for relatives only was presence of psychosis that did not meet DSM-III-R criteria for schizophrenia or schizoaffective disorder.

Clinical/Neuropsychological Assessment. All patients and relatives were interviewed using the Schedule for Affective Disorders and Schizophrenia–Lifetime Version (Spitzer and Endicott, 1978). The data collected was later supplemented with information from case-notes and from other relatives to assign a lifetime DSM-III-R psychiatric diagnosis. Structured assessments of schizotypal personality disorder were completed using the paranoid/schizoid and schizotypal personality subscale of the Family Interview for Genetic Studies (Nurnberger et al., 1994) derived from a reliable informant. Psychotic symptoms were assessed using PANSS (Kay et al., 1987) and SANS/SAPS (Andreasen, 1983, 1984). Overall functioning was assessed using GAF (American Psychiatric Association, 1994). Current IQ was estimated using 8 subtests of the Wechsler Adult Intelligence Scale–Revised (WAIS-R; Wechsler, 1981): Vocabulary, Comprehension, Information, Similarities, Block Design, Object Assembly, Picture Completion, and Picture Arrangement. Additionally, Vocabulary was used to estimate premorbid IQ. Processing speed was assessed using WAIS-R Digit Symbol and a computerized version of the Trail Making Test—Part A and B (Army Individual Test Battery, 1944; Reitan, 1958). Attention/vigilance was assessed using the CANTAB Rapid Visual Information Processing subtest. Verbal working memory was tested using the WAIS-R subtests Digit Span and Arithmetic. Spatial working memory was tested using the CANTAB Spatial Working Memory subtest (Cambridge Cognition, 2003a). Verbal and visual memory were assessed by the Wechsler Memory Scale–Revised (WMS-R) Logical Memory and Visual Reproduction subtests (Wechsler, 1987), respectively. Verbal and visual learning were measured using the WMS-R Paired Associate Learning and Visual Reproduction subtest, respectively.

MRI Acquisition. Not applicable.

Genotyping. DNA from a proportion of participants was obtained from whole blood or saliva and genotyped on the Affymetrix 6.0 array.

Ethics/Consent. The study was approved by the UK Multicenter Research Ethics Committee. All participants gave written informed consent before participating.

Acknowledgements. Support was provided by the European Community's Sixth Framework Programme through a Marie Curie Training Network (MRTN-CT-2006-035987) called the European Twin Study Network on Schizophrenia (EUTwinsS), NARSAD (through a Young Investigator Award to Dr. Touloupoulou), Wellcome Trust Research Training Fellowship (grant 064971 to Dr. Picchioni), Economic and Social Research Council/Medical Research Council and the Psychiatry Research Trust (PTA-037-27-0002); Deutsche Forschungsgemeinschaft (grant Et 31/2-1 to Dr. Ettinger), and Wellcome Trust Training Fellowship (ref 059007 to Dr. McDonald). The investigators acknowledge support from the Department of Health via the National Institute for Health Research (NIHR) Specialist Biomedical Research Centre for Mental Health award to South London and Maudsley NHS Foundation Trust (SLaM) and the Institute of Psychiatry at King's College London. Principal funding for genotyping was provided by the Wellcome Trust, as part of the Wellcome Trust Case Control Consortium 2 project (Grant Nos. 085475/B/08/Z and 085475/Z/08/Z). Dr. Bramon was supported by a MRC New Investigator Award, a MRC Centenary Award, the National Institute of Health Research UK (post-doctoral fellowship), the Psychiatry Research Trust, the Schizophrenia Research Fund, a Brain and Behavior Research Foundation (NARSAD) Young Investigator Award, a Wellcome Trust Research Training Fellowship, and the NIHR Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and Institute of Psychiatry Kings College London.

Conflicts of interest. Dr. Robin Murray has received honoraria for lectures from Janssen, AstraZeneca, Lilly, Novartis and Bristol-Myers Squibb. Dr. Picchioni has received travel awards from Pfizer, Janssen-Cilag, and Eli Lilly and an educational grant from Janssen-Cilag.

KCL-MTS — King's College London - Maudsley Twin Study

(Owens et al., 2012; Owens et al., 2011a; Owens et al., 2011b; Psychosis Endophenotypes International Consortium et al., 2014; Touloupoulou et al., 2007)

Recruitment. Probands with schizophrenia were recruited nationally throughout the United Kingdom from National Health Service treatment centers through referrals by their treating psychiatrist. In the United Kingdom, the National Health Service is a comprehensive national treatment service funded centrally and free at the point of delivery for all aspects of health care. By virtue of its financial and organizational structure, it is hugely inclusive and as a consequence cares for most patients with schizophrenia, making it a very representative system from which to recruit. Control twins were recruited from a volunteer twin register held at the Institute of Psychiatry, London, England. Zygosity was determined by assessment of 12 highly polymorphic microsatellite markers and a standardized twin likeness questionnaire. Medication status was recorded at the time of the assessment, and age at first contact with psychiatric services was ascertained to serve as a proxy index of age at illness onset. In concordant pairs, both members fulfilled criteria for DSM-IV schizophrenia or schizoaffective disorder. In discordant pairs, one member was diagnosed with DSM-IV schizophrenia, whereas the co-twin was free of any psychotic illness. In control pairs, both members were free of personal or family history of psychosis or schizophrenia spectrum disorder. Some participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. Exclusion criteria applied to all groups were: <18 years of age, a history of a neurological disorder or systemic illness with known neurological complications, a history of significant head injury associated with loss of consciousness for more than 1 minute, and current harmful substance use or dependence (defined as within the last 12 months). No participant included in the study had a psychotic illness directly attributable to the harmful use of illicit substances.

Clinical/Neuropsychological Assessment. The DSM-IV diagnoses were made using the Schedule for Affective Disorders and Schizophrenia (SADS)–Lifetime Version (Spitzer and Endicott, 1978), supplemented by information from medical notes or by using the Structured Clinical Interview for DSM-IV (First et al., 1995a, b). Psychotic symptoms in the probands in the month before testing were assessed using PANSS (Kay et al., 1987) and/or SANS/SAPS (Andreasen, 1983, 1984), and overall functioning using GAF (American Psychiatric Association, 1994). The full UK version of the WAIS-III (WAIS-III-UK; Wechsler, 2005) was administered to all participants to assess current IQ, consisting of the following subtests: Vocabulary, Comprehension, Information, Similarities, Letter-Number Sequencing, Digit Span, Arithmetic, Digit Symbol, Block Design, Object Assembly, Picture Completion, and Picture Arrangement. Additionally, Vocabulary was used to estimate premorbid IQ. Furthermore, processing speed was assessed using a computerized version of the Trail Making Test—Part A and B (Army Individual Test Battery, 1944; Reitan, 1958), Semantic Verbal Fluency (categories: animals, fruits, body parts) (Benton and Hamsher, 1989) and COWAT Phonemic Verbal Fluency (letters: F, A, S) (Reitan, 1985). Attention/vigilance was assessed using the CANTAB Rapid Visual Information Processing subtest (Cambridge Cognition, 2003a). Verbal working memory was tested using the WAIS-R subtests Letter-Number Sequencing, Digit Span and Arithmetic. Spatial working memory was tested using the WMS-R Visual Memory Span subtest (Wechsler, 1987) and the CANTAB Spatial Working Memory subtest (Cambridge Cognition, 2003a). Verbal and visual memory were assessed by the WMS-R Logical Memory and Visual Reproduction subtests (Wechsler, 1987), respectively. Verbal and visual learning were measured using the WMS-R Paired Associate

Learning subtest and Visual Reproduction/Visual Paired Associates subtests, respectively (Wechsler, 1987). Executive processing was assessed by the CANTAB Intra-Extra Dimensional Set Shifting subtest (Cambridge Cognition, 2003a).

MRI Acquisition. T1-weighted structural scans (TE = 5.0 ms; TI = 450 ms; TR = 35 ms; Flip angle = 30°; FOV = 20 cm; Spatial resolution = 0.78 x 0.78 x 1.5 mm; matrix = 256 x 256 x 128 mm; Slice acquisition direction = coronal) and Diffusion Tensor Imaging (TE = 102 ms; TR = 15000 ms; Flip angle = n/a; FOV = 24 cm; Spatial resolution = 2.5 x 2.5 x 2.5 mm; matrix = 96 x 96 mm; Slice acquisition direction = axial, AC-PC; number of diffusion-encoding gradient directions = 64; number of b0 volumes = 7; b-value = 1300 s/mm²) were acquired on a 1.5 Tesla GE Signa Advantage scanner using a circularly polarized (CP) head coil.

Genotyping. DNA from a proportion of participants was obtained from whole blood or saliva and genotyped on the Affymetrix 6.0 array.

Ethics/Consent. The study was approved by the UK Multicenter Research Ethics Committee. All participants gave written informed consent before participating.

Acknowledgements. Support was provided by the European Community's Sixth Framework Programme through a Marie Curie Training Network (MRTN-CT-2006-035987) called the European Twin Study Network on Schizophrenia (EUTwinsS), NARSAD (through a Young Investigator Award to Dr. Touloupoulou), Wellcome Trust Research Training Fellowship (grant 064971 to Dr. Picchioni), Economic and Social Research Council/Medical Research Council and the Psychiatry Research Trust (PTA-037-27-0002). The investigators acknowledge support from the Department of Health via the National Institute for Health Research (NIHR) Specialist Biomedical Research Centre for Mental Health award to South London and Maudsley NHS Foundation Trust (SLaM) and the Institute of Psychiatry at King's College London. Principal funding for genotyping was provided by the Wellcome Trust, as part of the Wellcome Trust Case Control Consortium 2 project (Grant Nos. 085475/B/08/Z and 085475/Z/08/Z). Dr. Bramon was supported by a MRC New Investigator Award, a MRC Centenary Award, the National Institute of Health Research UK (post-doctoral fellowship), the Psychiatry Research Trust, the Schizophrenia Research Fund, a Brain and Behavior Research Foundation (NARSAD) Young Investigator Award, a Wellcome Trust Research Training Fellowship, and the NIHR Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and Institute of Psychiatry Kings College London.

Conflicts of interest. Dr. Robin Murray has received honoraria for lectures from Janssen, AstraZeneca, Lilly, Novartis and Bristol-Myers Squibb. Dr. Picchioni has received travel awards from Pfizer, Janssen-Cilag, and Eli Lilly and an educational grant from Janssen-Cilag.

L&R — Language and Risk in Schizophrenia

(Francis et al., 2012)

Recruitment. Individuals at familial high risk for schizophrenia (FHR) (age range of 19–32 years) with at least one first degree family member suffering from schizophrenia or schizoaffective disorder and one second or third degree relative with history of a psychosis, suicide, or psychiatric hospitalization were included in this study. Participants were recruited during 2009–2011 from throughout Massachusetts and other New England neighboring regions through brochures and advertisements and by networking through the National Alliance on Mental Illness. Controls with no family history of a psychosis in 1st, 2nd or 3rd degree relatives (age range of 20–32 years) were also recruited from the community via advertisements.

Inclusion/Exclusion Criteria. Participants with a DSM-IV diagnosis of any history of lifetime psychotic disorder were excluded. Additional exclusion factors were: English not the participant's native language, non-right-handedness, neurological illness, and IQ below 80.

Clinical/Neuropsychological Assessment. All participants were administered the Diagnostic Interview for Genetic Studies (DIGS; Nurnberger et al., 1994). A family pedigree was drawn with information obtained from the participant. DSM-IV diagnoses (Axis I for major psychiatric illness

and Axis II for personality disorders) were made on all participants by a research psychiatrist using all available information collected during the interview. Schizotypal symptoms were measured using the Modified Structured Interview for Schizotypy (SIS; Kendler et al., 1989) from the DIGS. Data were acquired on 9 factors from the SIS (magical thinking, ideas of reference, illusions, suspiciousness, psychotic-like symptoms, restricted emotion, social isolation/introversion, schizotypal social anxiety, and anger to slights). Participants were administered a large neurocognitive test battery including tests estimating premorbid IQ, i.e. the Reading subtest from the WRAT-4 (Wilkinson and Robertson, 2006a), and current IQ, i.e. the Vocabulary and Block Design subtest of the WASI (Wechsler, 1999). BACS Symbol Coding (Keefe et al., 2008), Trail Making Test—Part A (Reitan, 1958), and Semantic Verbal Fluency (MCCB Category Fluency; category: animals) (Benton and Hamsher, 1978; Kern et al., 2008; Nuechterlein et al., 2008) were used for processing speed; the CPT-IP (Cornblatt et al., 1988) was used to measure attention/vigilance; the WMS-III Spatial Span subtest (Wechsler, 1997b) and University of Maryland Letter-Number Span (Gold et al., 1997) were used for assessing working memory. The ‘vigilance’, ‘working memory’ and ‘working memory plus interference’ measures from the Seidman Auditory CPT (Seidman et al., 1998; Seidman et al., 2006). The HVLT-R (Brandt and Benedict, 1997) and the WMS-III Logical Memory subtest (Wechsler, 1997b) assessed verbal learning and memory. The BVMT-R (Benedict, 1997) assessed visual learning. Executive function was measured using NAB Mazes (Stern and White, 2003) and the computerized WCST (Heaton, 2003; Heaton et al., 1993; Nestor et al., 1993). The D-KEFS Proverbs subtest (Delis et al., 2001; Delis et al., 2004) was administered to assess verbal ability, and Grooved Pegboard (Lafayette Instrument Company -- Trites, 1989) and Finger Tapping (Ruff and Parker, 1993) were administered to assess motor control. Additionally, participants were administered verbal neuropsychological tests to test many aspects of linguistic functioning potentially associated with familial high-risk for schizophrenia, including: Abstract Reasoning (Delis et al., 2001), Woodcock-Johnson Passage Comprehension Test (Woodcock et al., 2001); Phonological Awareness derived from auditory stimuli: Elision and Blending Words subtests from the Comprehensive Test of Phonological Processing (Wagner et al., 1999); Phonological Awareness derived from visual stimuli including real words (WRAT-4 Reading subtest; Test of Word Reading Efficiency [TOWRE] Sight Word Efficiency; Woodcock-Johnson Letter Word Identification) and non-words (TOWRE Phonemic Decoding Efficiency; Woodcock-Johnson Word Attack) (Torgesen et al., 1999; Wilkinson and Robertson, 2006b; Woodcock et al., 2001).

MRI Acquisition. T1-weighted structural scans (TE = 3.48 ms; TI = 900 ms; TR = 15 ms; Flip angle = 9°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.0 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 84 ms; TR = 8060 ms; Flip angle = 55°; FOV = 25.6 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 60; number of b0 volumes = 10; b-value = 700 s/mm²) on a 3 Tesla Siemens Trio Tim scanner using a 32-channel head coil.

Genotyping. DNA was obtained from whole blood and genotyped on the Illumina Infinium PsychArray. Genotyping was funded by the National Institute of Mental Health (NIMH) of the National Institutes of Health (NIH) grant number R01MH092380 to T.L.P. supporting the GENUS Consortium.

Ethics/Consent. The study was approved by the Human Subjects Investigation Committee at Harvard Medical School, Beth Israel Deaconess Medical Center, Massachusetts Institute of Technology, Brigham and Women's Hospital, and Veterans Administration Boston Healthcare System, Brockton, Massachusetts. All participants provided written informed consent and were paid for their participation.

Acknowledgements. Data acquisition was funded by NIMH grant R21MH083205 (LE DeLisi), R01MH064023 (MS Keshavan), and the Commonwealth Research Center of the Massachusetts Department of Mental Health grant SCDMH82101008006 (LJ Seidman).

MCIC — Mind Clinical Imaging Consortium

(Gollub et al., 2013)

Recruitment. Patients with schizophrenia and demographic, age and sex- matched healthy control participants were recruited for this study between July 2004 and July 2006 across four research sites—University of New Mexico, University of Minnesota, Massachusetts General Hospital, and University of Iowa. The final cohort for whom cognitive and/or MRI data are available includes 156 patients and 122 controls. However, the University of Iowa subsample (38 patients, 25 controls) is not included in the GENUS dataset, resulting in 118 patients and 97 controls included in the GENUS dataset. Participants form part of the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Schizophrenia Working Group dataset (van Erp et al., 2016).

Inclusion/Exclusion Criteria. Participants were excluded for reasons such as excessive motion during image acquisition, claustrophobia or discomfort during MRI scanning, or a determination of ineligibility based on screening exams and tests. All participants were between the ages of 18 and 60 and spoke English as their native language. To be included in the schizophrenia cohort, patients had to meet DSM-IV diagnostic criteria for schizophrenia, schizoaffective disorder or schizophreniform disorder. Concerted effort was made to recruit patients early in the course of their illness and especially those who were antipsychotic drug naïve. The healthy control participants with no current or past history of psychiatric illness including substance abuse or dependence were matched within site to the patient cohort for age, sex, and parental education. Control participants who had not been diagnosed with any psychiatric disorders, but had been medicated with antidepressants, anti-anxiety medication or medication for sleep disturbance were included in the study, provided that the duration of their medication did not exceed 2 months of lifetime use and no medication was used within the 6 months preceding the baseline MRI scan. Control participants who met criteria for current or past history of substance abuse or dependence were excluded from the study. Patients were not excluded from the study unless criteria were met for current (i.e., within the past month) abuse or dependence except for 6 patients who were found to meet criteria for current abuse after the study data was collected. Both patients and controls were excluded if they had 1) an IQ less than 70 based on a standardized IQ test, 2) history of a head injury resulting in prolonged loss of consciousness, neurosurgical procedure, neurological disease, history of skull fracture, severe or disabling medical conditions, or 3) a contraindication for MRI scanning such as pregnancy, metal in body or head including implanted pacemaker, medication pump, vagal stimulator, deep brain stimulator, implanted TENS unit, or ventriculo-peritoneal shunt. Clinical teams at each site recruited the patients. Primary treating clinicians, not necessarily involved with the study, identified potential patients who met inclusion/ exclusion criteria and were competent to give informed consent. Healthy control participants were recruited from the community at each site through posters, brochures, newspapers, website advertisements, and institutional subject recruitment resources.

Clinical/Neuropsychological Assessment. Core demographic information collected includes age, handedness, height, weight, ethnicity, race, educational achievement, parental education, and socioeconomic status. A Structured Clinical Interview for DSM-IV Disorders, SCID-P for patients and SCID-NP for controls (First et al., 1995a, b), or the Comprehensive Assessment of Symptoms and History (CASH; Andreasen et al., 1992) were used to diagnose primary and co-morbid psychiatric disorders in controls and patients. Additionally, patients were interviewed with the SANS/SAPS (Andreasen, 1983, 1984) and the CDSS (Addington et al., 1992) to record symptoms and their current severity. Premorbid IQ and current IQ were measured using the Reading subtest of the WRAT-3 (Wilkinson, 1993), and the Block Design, Vocabulary, and Similarities subtests of the WAIS-III (Wechsler, 1997a), respectively. Trail Making Test—Part A and B (Reitan, 1958) measured processing speed. D-KEFS (Delis et al., 2001) assessed Semantic Verbal Fluency (categories: animals, fruits) and Phonemic Verbal Fluency (letters: F, A, S). The WAIS-III Letter-Number Sequencing subtest measured working memory. Verbal

memory and learning were assessed using the Wechsler Memory Scale—Third Edition (WMS-III) Logical Memory subtest (Wechsler, 1997b) and the HVLT-R (Brandt, 1991). The Benton Visual Retention Test (BVRT; Sivan, 1992) and the Face Recognition subtest of the WMS-III (Wechsler, 1997b) both measured visual memory. A computerized version of the Tower of London test (TOL; Shallice, 1982) was employed to measure planning and problem solving. The Grooved Pegboard Test (Ruff and Parker, 1993), and the California Computerized Assessment Package (CalCap; Miller, 1990), a computerized reaction time test, tested fine motor dexterity and speed. The Annett Scale of Hand Preference (Annett, 1970) provided a numeric measure of handedness.

MRI Acquisition. T1-weighted structural scans at Massachusetts General Hospital, Martinos Center (TE = 4.76 ms; TI = 300 ms; TR = 12.0 ms; Flip angle = 20°; FOV = 16 cm; Spatial resolution = 0.7 x 0.7 x 1.5 mm; matrix = 256 x 256 x 128 mm; Slice acquisition direction = axial, oblique) and Diffusion Tensor Imaging (TE = 80 ms; TR = 8900 ms; Flip angle = 90°; FOV = 16 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 60; number of b0 volumes = 10; b-value = 700 s/mm²) were acquired on a 1.5 Tesla Siemens Sonata scanner using an 8-channel head coil. T1-weighted structural scans at the University of Minnesota (TE = 3.81 ms; TI = 1100 ms; TR = 2530 ms; Flip angle = 7°; FOV = 16 cm; Spatial resolution = 0.625 x 0.625 x 1.5 mm; matrix = 256 x 256 x 128 mm; Slice acquisition direction = coronal) and Diffusion Tensor Imaging (TE = 98 ms; TR = 11500 ms; Flip angle = 90°; FOV = 16 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 12; number of b0 volumes = 1; b-value = 1000 s/mm²) were acquired on a 3.0 Tesla Siemens Trio Tim scanner using an 8-channel head coil. T1-weighted structural scans at the University of New Mexico (TE = 4.76 ms; TI = 300 ms; TR = 12.0 ms; Flip angle = 20°; FOV = 16 cm; Spatial resolution = 0.625 x 0.625 x 1.5 mm; matrix = 256 x 256 x 128 mm; Slice acquisition direction = coronal) and Diffusion Tensor Imaging (TE = 92 ms; TR = 10000 ms; Flip angle = 90°; FOV = 16 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 12; number of b0 volumes = 1; b-value = 1000 s/mm²) were acquired on a 1.5 Tesla Siemens Sonata scanner using an 8-channel head coil.

Phenotype data are available from the Collaborative Informatics and Neuroimaging Suite (<https://coins.mrn.org>).

Genotyping. DNA was obtained from whole blood and genotyped on the Illumina HumanOmni1-Quad BeadChip.

Ethics/Consent. All participants provided informed consent to participate in the study. The study was approved by the human research committees at each site. In addition to informed consent, all patients successfully completed a questionnaire verifying that they understood the study procedures.

Acknowledgements. The Mind Clinical Imaging Consortium was supported primarily by the Department of Energy DE-FG02-99ER62764 through its support of the Mind Research Network (MRN, formerly known as the MIND Institute) and the consortium, as well as by a NARSAD Young Investigator Award (to Dr. Ehrlich), the Blowitz-Ridgeway and Essel Foundations, NWO ZonMw TOP 91211021, DFG research fellowship (to Dr. Ehrlich), the Mind Research Network, National Institutes of Health through NCRR 5MO1-RR001066 (MGH General Clinical Research Center), NIMH K08 MH068540, the Biomedical Informatics Research Network with NCRR Supplements to P41 RR14075 (MGH), M01 RR 01066 (MGH), NIBIB R01EB006841 (MRN), R01EB005846 (MRN), 2R01 EB000840 (MRN), 1RC1MH089257 (MRN), as well as grant U24 RR021992, P20RR021938/P20GM103472 and R01MH094524. The investigators wish to thank their many colleagues who served as mentors, advisors and supporters during the inception and conduct of the study including Donald Goff, Gina Kuperberg, Jill Goldstein, Martha Shenton, Robert McCarley, Stephan Heckers, Cynthia Wible, Raquelle Meshulam-Gately, and Mark Vangel. The investigators thank the study staff and clinicians at each site that were responsible for the data

acquisition, including Stuart Wallace, Ann Cousins, Raquelle Mesholam-Gately, Steven Stufflebeam, Oliver Freudenreich, Daphne Holt, Joshua Roffman, Laura Kunkel, Frank Fleming, George He, Hans Johnson, Ron Pierson, Arvind Caprihan, Phyllis Somers, Christine Portal, Kaila Norman, Diana South, Michael Doty and Haley Milner. The investigators also acknowledge the expert guidance on image and other types of data acquisition obtained from Lee Friedman, Stephan Posse, Jorge Jovicich, and Tom Wassink, and acknowledge the many research assistants, students and colleagues who assisted in data curation over the years since data acquisition was completed, including Stuart Wallace, Carolyn Zyloney, Komal Sawlani, Esther Walton, Jill Fries, Adam Scott, Dylan Wood, Runtang Wang, William Courtney, Angie Guimaraes, Lisa Shenkman, Mustafa Kendi, Aysa Tuba Karagulle Kendi, Ryan Muetzel, Tara Biehl, and Marcus Schmidt.

MGH — Massachusetts General Hospital

(Ho et al., 2016; Holt et al., 2012; Linnman et al., 2013; Roffman et al., 2013; Roffman et al., 2011; Roffman et al., 2007; Roffman et al., 2008a; Roffman et al., 2008b)

Recruitment. Outpatients with schizophrenia were recruited from an urban community mental health center in Boston. Demographically matched healthy participants were recruited from the community by poster and website advertisements.

Inclusion/Exclusion Criteria. Participants were excluded if they had a history of substance abuse or dependence within the previous 6 months, a history of significant head injury, or neurologic illness. Healthy participants were screened to exclude a personal history of Axis I mental illness (First et al., 2002a) or a family history of schizophrenia-spectrum disorder. For all participants, exclusion criteria included severe medical illness, significant head trauma, neurologic illness, substance abuse during the past six months and contraindications for MRI scanning (e.g., implanted metal objects, claustrophobia, renal insufficiency). Patients had been maintained on stable doses of second generation antipsychotics for at least six weeks, with the exception of one patient taking prolixin, and three patients who were not taking any medications.

Clinical/Neuropsychological Assessment. Patient diagnoses were confirmed using the Structured Clinical Interview for DSM-IV-TR (First et al., 2002b). The healthy controls were without psychiatric disorders as determined by the Structured Clinical Interview for DSM-IV Axis I Disorders (First et al., 1995b). Participants were also rated with the PANSS (Kay et al., 1987), Hamilton Depression Rating Scale (HDRS; Hamilton, 1960) and Simpson–Angus Scale (Simpson and Angus, 1970), a measure of extrapyramidal symptoms. Tests were administered by one of four trained and certified raters. Premorbid IQ was estimated using the North American Adult Reading Test (NAART; Blair and Spreen, 1989). The WAIS—III (Wechsler, 1997a) was conducted to measure current IQ (all subtests [=13], except Symbol Search). Patients participated in a neurocognitive battery that included the CVLT (Delis et al., 1987), Semantic Verbal Fluency (category: animals) (Benton and Hamsher, 1989) and COWAT Phonemic Verbal Fluency (letters: F, A, S) (Reitan, 1985), WCST (Heaton et al., 1993), and Finger Tapping (Ruff and Parker, 1993).

MRI Acquisition. Subsample1: T1-weighted structural scans (TE = 1.54 ms, 3.36 ms; TI = 1100 ms; TR = 2200; Flip angle = 7°; FOV = 23 cm; Spatial resolution = 1.2 x 1.2 x 1.2 mm; matrix = 192 x 192 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 85.0 ms; TR = 5980 ms; Flip angle = 90°; FOV = 22 cm; Spatial resolution = 1.375 x 1.375 x 3.0 mm; matrix = 160 x 160 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 6; number of b0 volumes = 1; b-value = 1000 s/mm²) were acquired on a 3 Tesla Siemens Trio Tim scanner. Subsample 2: T1-weighted structural scans (TE = 3.39 ms; TI = 1100 ms; TR = 2530 ms; Flip angle = 7°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.3 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) were acquired on a 3 Tesla Siemens Trio Tim scanner using a 12-channel head coil.

Genotyping. DNA was obtained from whole blood or saliva and genotyped on the Illumina Infinium PsychArray. Genotyping was funded by the National Institute of Mental Health (NIMH) of

the National Institutes of Health (NIH) grant number R01MH092380 to T.L.P. supporting the GENUS Consortium.

Ethics/Consent. Study procedures were approved by the institutional review boards of Partners HealthCare and the Massachusetts Department of Mental Health. All participants provided written informed consent.

Acknowledgements. The collection of the cognitive and clinical subsample was supported by NIH grants MH02025-01A3, MH60450 (Dr. Goff) and DK56085 (Dr. Halsted), as well as by a NARSAD Independent Investigator Award and the Sidney R. Baer Jr. Foundation (Dr. Goff), and an APIRE/Lilly Psychiatric Research Fellowship and Harvard Medical School Dupont-Warren Fellowship (Dr. Roffman). The acquisition of neuroimaging data for subsample 1 was funded by National Institute of Mental Health grants R01MH067720 (Dr. Manoach), R01MH070831 (Dr. Goff), R01MH101425 and K23MH084059 (Dr. Roffman), a grant from PamLab (Dr. Roffman), and the Howard Hughes Medical Institute Physician-Scientist Early Career Award (Dr. Roffman). This work was conducted with support from Harvard Catalyst, The Harvard Clinical and Translational Science Center (National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health award UL1 RR 025758, and financial contributions from Harvard University and its affiliated academic health care centers). The acquisition of neuroimaging data for subsample 2 was supported by National Institute of Mental Health grants K23MH076054 and R01MH095904 (Dr. Holt) and NARSAD with the Sidney R. Baer, Jr. Foundation (Dr. Holt). The investigators thank David C. Henderson, Alexandra S. Tanner, Madeline Giegold, and Noah J. Silverstein, as well as Garth Coombs III, Adam Z. Nitenson, and Liana J. Petrucci for their indispensable help with the MRI and clinical data acquisition, and Stephanie N. DeCross and Franklin C. Huntington for administrative support.

NEFS — New England Family Study

(Buka et al., 2013; Goldstein et al., 2010; Goldstein et al., 2014; Seidman et al., 2013; Seidman et al., 2002)

Case and Control Recruitment. Participants were selected from the Boston and Providence cohorts of the Collaborative Perinatal Project (CPP), currently known as the New England Family Studies (NEFS). The CPP was initiated over 50 years ago to investigate prospectively the prenatal and familial antecedents of pediatric, neurological and psychological disorders of childhood (Niswander and Gordon, 1972). Approximately 17,000 pregnancies were followed in New England. Many types of assessments, including psychological examinations, were conducted on offspring up to 7 years of age, when the study officially ended in 1973. Ascertainment and assessment of the adult psychotic patients in NEFS were identified through a systemic follow-up of the entire NEFS cohort of the CPP. NEFS parents and offspring with a history of psychiatric hospitalization and/or possible psychotic and bipolar illness were identified from the following sources: (1) record linkages with public hospitals, mental health clinics, and the Massachusetts and Rhode Island Departments of Mental Health; (2) several follow-up and case-control studies nested within the larger NEFS cohort, involving direct interviews with approximately 20% of the cohort; and (3) reports from participants in these interview studies of a family member with a history of psychotic or bipolar symptoms or diagnosis. Controls were selected from families participating in the control arm of a NEFS high-risk study in which the control population was a random stratified sample of parents selected from the entire NEFS cohort, with no known history of psychosis or other major Axis I disorders. Thus, the NEFS sample of cases and controls is a representative community sample of subjects.

Inclusion/Exclusion Criteria. Adult offspring with major non-organic psychoses (including schizophrenia, schizoaffective disorder, delusional disorder, brief psychosis, psychosis NOS, bipolar disorder with psychotic features, and major depressive disorder with psychosis) within NEFS cohorts were identified approximately 30 years later through a two-stage diagnostic assessment procedure between 1996 and 2007. The investigators were blind to prior

assessments of these subjects during this follow-up. In the first stage, individuals with possible psychotic illness were identified through record linkages and from personal interviews. In the second stage, those who consented to participate in follow-up efforts were interviewed using the SCID (First et al., 1995b). Based on interview data and medical record review, trained PhD- and MD-level diagnosticians then completed best-estimate consensus diagnoses according to DSM-IV criteria for lifetime prevalence of psychotic and other psychiatric disorders. Exclusion criteria for all adult participants were a history of neurological disease, traumatic brain injury, medical illness or alcohol-related disease with documented cognitive sequelae, major sensory impairments (e.g. deafness), IQ<65 in adulthood or inability to understand the procedures, <6 years of formal education and severe substance abuse within the past 6 months. All psychoses cases were living in the community when assessed. Controls had to be free of any known lifetime history of psychosis or other major Axis I disorders. Furthermore, parents and grandparents, in addition to the parents' siblings, had to be free of any known lifetime history of psychosis, bipolar, schizotypal, recurrent MDD, suicide attempts or psychiatric hospitalizations. Siblings of the controls also had to be free of any lifetime history of psychosis or BD.

Family High Risk Sample Ascertainment: The study goal was to ascertain approximately 200 psychotic NEFS Generation-1 parents, half with schizophrenia and half with affective psychoses and a comparable group of normal control parents. Generation-1 parents were contacted between 1994 and 2002. Parents with a history of psychiatric treatment were identified by the following sources: (1) prior record review; (2) subsequent record linkages with private and public psychiatric treatment facilities in Rhode Island and Massachusetts; (3) information provided by mothers during the original study (e.g., psychiatric hospitalization at study enrollment; psychiatric treatment at the 7 year assessment; or history of treatment with antipsychotic medication); or (4) family member reports from recent follow-up studies with the NCPP cohort. Through these efforts, from a total pool of 26,928 G1 parents, we identified 859 persons with indications of potential psychosis. This represents a rate of approximately 3.2%-4.2% of the pool of parents we were able to review, which is consistent with current estimates of the lifetime prevalence of psychotic disorders. Of these parents, 755 were eligible for follow-up. Parents were considered eligible if at least one Generation-2 offspring was assessed after 4 months of age.

Control parents were selected to be comparable to parents with psychotic disorders based on the number of offspring enrolled in the CPP, patient status (public or private), parent's age, ethnicity (Caucasian or other), study site, and on the offspring's age, sex, and history of chronic hypoxia (given that we wanted to test for the interaction of genetic vulnerability and this obstetric condition). Eligible controls included all members of the CPP who were not identified as potential psychotic parents and whose records did not indicate a history of psychiatric treatment. Healthy control parents did not have spouses, parents, siblings, or any second-degree relatives with psychoses, recurrent MDD, suicide, or psychiatric hospitalizations.

Generation-1 Follow-Up and Psychiatric Assessment: Subjects were located through a variety of methods, including thorough searches of credit bureaus, address directories, death certificates, motor vehicle reports, and by home visits. The located subjects were invited to participate in a two-part interview. The first interview screened for potential psychoses and the second was a full diagnostic interview using the SCID (First et al., 1995b) and assessing Axis I diagnoses of any form of psychotic, major affective, and bipolar disorders; and substance abuse or dependence (alcohol and drug). The clinical interview was conducted by systematically trained M.A.-level clinical interviewers. Medical records were obtained with subject consent. Family history of psychiatric disorders was evaluated using the Family Interview for Genetic Studies (Maxwell, 1996). Expert diagnosticians (including J.G., L.S.) reviewed all of the information collected from both interviews and medical records, if available, to determine final best estimate diagnoses. If there was any evidence of mental illness in the spouse, we interviewed the spouse or searched for medical records if the spouse had passed away. Spouses' names were subjected to the linkage procedures used for the mothers.

Of the 755 eligible Generation-1 parents, 212 with confirmed DSM-IV psychotic disorders were identified, with 153 (72%) mothers and 59 (27%) fathers. The sample of 212 parents with psychosis and 132 non-psychotic parents had a total of 467 pregnancies: 167 offspring among affective parents, 114 offspring among SPS parents, and 186 offspring among non-psychotic parents. Of note, among the 344 final Generation-1 parents, there are six 'two parent families' in which both parents were diagnosed with some form of psychosis, resulting in a sample size of 338 unduplicated families. Of the 467 pregnancies, 22 (4.7%) resulted in an offspring who was stillborn, did not live past early childhood, was lost to follow-up before age 7, or was adopted/placed in foster care, and thus ineligible for this adult follow-up study. Of the 445 eligible Generation-2's, 428 (96.2%) were located of whom 344 (80.4%) were interviewed, for an overall interview rate of 77.3%. In addition to the 344 Generation-2's with diagnostic information from interviews, there were an additional six Generation-2's with a diagnostic summary from medical records from treatment facilities for a total of 350 Generation-2's with diagnostic information. This reflects a completed diagnostic rate of 78.7%.

Clinical/Neuropsychological Assessment. Trained and skilled M.A.-level interviewers and neuropsychological testers conducted all clinical and cognitive assessments. The SANS/SAPS (Andreasen, 1983, 1984)) were used to measure symptom severity. Participants were administered a large neurocognitive test battery including tests estimating premorbid IQ (the Reading subtest from the WRAT-3 (Wilkinson, 1993), and current IQ (the Vocabulary, Information, Comprehension, Block Design, Picture Arrangement, Digit Span and Digit Symbol subtests of the WAIS-R (Wechsler, 1974). The WAIS-R Digit Symbol Coding subtest (Wechsler, 1974) and COWAT Phonemic Verbal Fluency test (letters F, A, S) (Reitan, 1985) were used for measuring processing speed. The 'vigilance', 'working memory' and 'working memory plus interference' measures from the Seidman Auditory CPT (Seidman et al., 1998; Seidman et al., 2006) were used to measure attention/vigilance and verbal working memory, in addition to the WAIS-R Digit Span subtest for working memory assessment (Wechsler, 1974). The CVLT or CVLT-II (Delis et al., 1987, 2000) and the WMS-R or WMS-III Logical Memory subtest (Wechsler, 1987, 1997b) assessed verbal learning and memory. The WMS-III Faces subtest (Wechsler, 1997b) and the Rey-Osterrieth Complex Figure Test (Rey, 1964) assessed visual learning and memory. Executive function was measured using the WCST (Heaton et al., 1993) or the Stroop test (Stroop, 1935). Finally, the Rapid Automatized Naming test (Denckla and Rudel, 1974) was used to measure language ability.

MRI Acquisition. T1-weighted structural scans (TE = 3.31 ms; TI = 1000 ms; TR = 2730 ms; Flip angle = 7°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.33 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 77-79 ms; TR = 7200 ms; Flip angle = 90°; FOV = 25.6 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 60; number of b0 volumes = 10; b-value = 700 s/mm²) were acquired on a 1.5 Tesla Siemens Avanto scanner using a 4-channel head coil. For a second subsample, T1-weighted structural scans (TE = 3.39 ms; TI = 1000 ms; TR = 2730 ms; Flip angle = 7°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.33 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 67 ms; TR = 9000 ms; Flip angle = 90°; FOV = 25.6 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 6; number of b0 volumes = 1; b-value = 600 s/mm²) were acquired on a 1.5 Tesla Siemens Sonata scanner using a CP coil. A portion of T1-weighted structural scans were acquired also on a 1.5 Tesla Siemens Sonata scanner (TE = 4.3 ms; TI = 8 ms; TR = 11.8 ms; Flip angle = 8°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.5 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal, CP head coil) or on a 1.5T GE Genesis Signa scanner (TE = 1.6 ms; TI = 300 ms; TR = 6.7 ms; Flip angle = 25°; FOV = 24 cm; Spatial resolution = 0.94 x 0.94 x 1.5 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal, CP head coil). Given that NEFS is a long-term follow up study, in more recent years, T1-

weighted scans were acquired on a 3 Tesla Siemens Trio Tim scanner, using a 12-channel head coil (TE = 3.39 ms; TI = 1000 ms; TR = 2530 ms; Flip angle = 7°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.33 mm; matrix = 256 x 256 mm; Slice acquisition direction = sagittal) and Diffusion Tensor Imaging (TE = 83-84 ms, TR = 9400-9440 ms; Flip angle = 90°; FOV = 25.6 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 60; number of b0 volumes = 10; b-value = 700 s/mm²).

Genotyping. DNA was obtained from whole blood or buffy coat and genotyped on the Illumina Infinium PsychArray. Genotyping was funded by the National Institute of Mental Health (NIMH) of the National Institutes of Health (NIH) grant number R01MH092380 to T.L.P. supporting the GENUS Consortium.

Ethics/Consent. Human subject research approval was granted by Institutional Review Boards at Harvard University, Partners Healthcare, Brown University and local psychiatric facilities. Written consent was obtained from all participants interviewed. Participants were compensated for their participation.

Acknowledgements. Original sample ascertainment was supported by NIMH RO1 MH50647 (1999–2003, Tsuang, P.I.; 2003–2006, Goldstein, P.I.), RO1 MH56956 (Goldstein, P.I.), and the Stanley Medical Research Institute (Buka and Goldstein). The authors would like to express their appreciation to the dedicated NEFS research team including Jasmina Burdzovic-Andreas, Ph.D., Monica Landi, M.S.W., June Wolf, Ph.D., and JoAnn Donatelli, Ph.D., for their contributions to the recruitment of subjects, clinical interviewing and diagnoses, Christiana Provencal, M.A., for her contribution to project management during the original sample acquisition. We also appreciate the efforts of a number of research assistants over the years and a previous project coordinator, Lisa Denny, MD, for their contributions to the study at an earlier stage of the work and Ronald O. Rieder, MD, for access to his earlier 1975 study of some of the Boston mothers in our parental case group. We would also like to thank Harlyn Aizley, Ed.M., Jenn Walch, M.Ed., for their immense contributions to the collection and management of data, respectively. The investigators furthermore wish to acknowledge Anne Remington, Lynda Tucker, and Dr. George Papadimitriou for their contributions.

PAGES — Phenomics and Genomics Sample

(Walters et al., 2010; Walters et al., 2013)

Recruitment. Clinically stable patients with a DSM-IV diagnosis of schizophrenia were ascertained from mental health services in the Munich area. Healthy control participants randomly selected from population registers from the Munich area were contacted by mail and invited to participate. The address, sex, and age of potential control participants were supplied by the municipal registration office, which enabled equal numbers of males and females and people of different ages (in decades) to be contacted. In patients, 68% were of German descent (all 4 grandparents born in Germany), and 32% were German white. All healthy control participants were of German descent (all 4 grandparents German). All patients and healthy control participants were unrelated. Some patients (62%) and healthy control participants (23%) form part of the SGENE-plus study (Stefansson et al., 2009). Some participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014). Controls form part of the Cognitive Genomics Consortium wave 1 (COGENT1) dataset (Lencz et al., 2014).

Inclusion/Exclusion Criteria. The inclusion criteria for patients were a diagnosis of schizophrenia (6-month symptom duration) and 18 to 80 years of age. Exclusion criteria were a history of head injury or neurological disease. Healthy control participants 18 to 65 years of age underwent an extensive screening process (Walters et al., 2010) to exclude those with neurological or psychotic disorders, those who had first-degree relatives with psychotic disorders, and those with lifetime exposure to psychiatric medication. In the case of healthy controls older than 60 years, the Mini-Mental State Examination (Folstein et al., 1983) was employed to exclude individuals with possible cognitive impairment. A neurologic examination was also conducted to exclude individuals with current central nervous system impairment.

Clinical/Neuropsychological Assessment. Detailed medical and psychiatric histories were collected, including data from a clinical interview using the SCID (First et al., 1995b) to evaluate lifetime Axis I and II diagnoses. The interviews were rated by four psychiatrists or one psychologist, and all measurements were independently rated by a senior researcher. Participants were also rated for symptoms using the PANSS (Kay et al., 1987). In addition, the Family History Assessment Module (Rice et al., 1995) was conducted to exclude psychotic disorders in first-degree relatives. Current IQ was measured by the full German WAIS-R (Tewes, 1991), consisting of 11 subtests: Vocabulary, Comprehension, Information, Digit Span, Arithmetic, Similarities, Block Design, Picture Completion, Picture Arrangement, Object Assembly, and Digit Symbol Coding. Additionally, Vocabulary was used to estimate premorbid IQ. All control participants completed the full WAIS-R assessment. All patients and 25% of control participants completed an extensive neuropsychological battery consisting of the WAIS-R and the following: Trail Making Test—Part A (Reitan, 1958), Semantic Verbal Fluency (categories: animals, food) (Benton and Hamsher, 1978), and Phonemic Verbal Fluency (letters: S and P) (Reitan, 1985) to assess attention/processing speed; Continuous Performance Test 3–7 Version (Nuechterlein and Asarnow, 2004) to assess attention/vigilance; WMS-R Spatial Span subtest (Härting et al., 2000; Wechsler, 1987) and a spatial n-back task (Callicott et al., 1998) to assess spatial working memory; Verbal Learning and Memory Test (Helmstaedter et al., 2001) or Verbaler Lerntest (Sturm and Willmes, 1999), and WMS-R Verbal Paired Associates subtest (Härting et al., 2000; Wechsler, 1987) to assess verbal learning and memory; WMS-R Logical Memory subtest (Härting et al., 2000; Wechsler, 1987) to assess episodic memory; WMS-R Figural Memory, Visual Reproduction, and Visual Paired Associates subtests (Härting et al., 2000; Wechsler, 1987) to assess visual learning and memory; and Trail Making Test-Part B (Reitan, 1958) and WCST (Heaton, 2003; Heaton et al., 1993) to assess executive function.

MRI Acquisition. Not applicable.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on the Illumina OmniExpress array or Illumina HumanHap300 array.

Ethics/Consent All participants provided written informed consent in accordance with the relevant ethics approvals.

Acknowledgements. Parts of the PAGES study were supported by the German Federal Ministry of Education and Research (BMBF) through the Integrated Network IntegraMent (Integrated Understanding of Causes and Mechanisms in Mental Disorders), under the auspices of the e:Med Programme (grant 01ZX1314).

PHRS — Pittsburgh High Risk Study

(Bhojraj et al., 2011; Bhojraj et al., 2009)

Recruitment. The study was conducted at the Western Psychiatric Institute and Clinic, Pittsburgh. First- or second-degree relatives of individuals diagnosed with schizophrenia and age- and gender-matched healthy comparison participants from an ongoing study at the University of Pittsburgh participated in this study. Relatives of patients with schizophrenia or schizoaffective disorder were recruited by approaching patients in the clinic and through advertisements. Controls were recruited through advertisements in the same community. The relatives and controls were included in the GENUS FHR and control groups, respectively.

Inclusion/Exclusion Criteria. Participants with an IQ < 80, lifetime evidence of a psychotic disorder, exposure to antipsychotic medications or anti-depressant medications, substance use disorder within the last month, significant neurological or medical condition were excluded.

Clinical/Neuropsychological Assessment. Clinical assessments of all participants, and parental diagnoses of schizophrenia or schizoaffective disorder, used the SCID (First et al., 1995b) and were confirmed using consensus meetings led by senior diagnosticians. None of the relatives were diagnosed with psychotic or other psychiatric illnesses and none had received antipsychotic medications. In addition to the SCID, the Structured Interview for Prodromal Syndromes, including the Scale of Prodromal Symptoms, was administered (Miller et al., 2003). Current IQ was assessed using the Block Design, Matrix Reasoning, Vocabulary, and Similarities subtests from the WASI (Wechsler, 1999). Verbal fluency was assessed using the Multilingual Aphasia Examination (Benton and Hamsher, 1978) including a Phonemic Verbal Fluency task (number of words generated in 20 seconds (each) that start with the letter C, F, and L) (Reitan, 1985) and a Semantic Verbal Fluency task (categories: animals, fruits and vegetables) (Benton and Hamsher, 1978). The CPT-IP (Cornblatt et al., 1988), AX-CPT, Spatial Working Memory, Word List Memory, and Go-No-Go subtests from Cogtest (Sharma and Bilder, 2004; The Cognition Group, 2006) were used to assess attention/vigilance, attention, spatial working memory, verbal learning and memory, and executive functioning, respectively. The Penn Visual Object Learning Test (Glahn et al., 1997), the Computerized Finger Tapping Test (Coleman et al., 1997), and the Penn Emotion Recognition Test (Gur et al., 2002) from the Computerized Neurocognitive Battery (CNB; (Gur et al., 2010; Moore et al., 2015; Swagerman et al., 2016)) were used to assess visual learning, motor control, and social cognition, respectively. Further, the WCST (Heaton et al., 1993) was used to assess executive function.

MRI Acquisition. T1-weighted structural scans (TE = 5.0 ms; TR = 25 ms; Flip angle = 40°; FOV = 24 cm; Spatial resolution = 0.94 x 0.94 x 1.5 mm; matrix = 256 x 192 mm; Slice acquisition direction = coronal, AC-PC oblique) were acquired on a 1.5 Tesla GE Genesis Signa scanner using a circularly polarized head coil.

Genotyping. DNA was obtained from whole blood and genotyped on the Illumina Infinium PsychArray. Genotyping was funded by the National Institute of Mental Health (NIMH) of the National Institutes of Health (NIH) grant number R01MH092380 to T.L.P. supporting the GENUS Consortium.

Ethics/Consent. The study was approved by the University of Pittsburgh Institutional Review Board. All participants signed informed consent after the study was fully explained to them. For participants <18 years of age, the consent was provided by the parent or guardian, and the participants provided informed assent.

Acknowledgements. The study was funded by NIMH grant MH64023 (Dr. Keshavan). The investigators acknowledge the contributions of Dr. Vishwajit L. Nimgaonkar, Dr. Kodavali Chowdari, Dr. Debra M. Mostrose, Jean M. Miewald, Ian Mathew, Brandon M. Hager, and Olivia Lutz.

TCD/NUIG — Trinity College Dublin and National University of Ireland, Galway

(Donohoe et al., 2009; O'Donovan et al., 2008; Walters et al., 2010; Walters et al., 2013)

Recruitment. Patients with schizophrenia were recruited from across Ireland. Comparison participants were recruited by local media advertisements. Some participants (~50%) form part of the Psychiatric Genomics Consortium Schizophrenia Working Group wave 2 dataset (PGC Schizophrenia Working Group, 2014). Some controls (~150) form part of the Cognitive Genomics Consortium wave 2 (COGENT2) dataset (Trampush et al., 2017). Some participants (~20%) form part of the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Schizophrenia Working Group dataset (van Erp et al., 2016).

Inclusion/Exclusion Criteria. Exclusion criteria were a history of head injury or neurological disease. The inclusion criteria required that participants had no history of substance abuse in the preceding 6 months based on self-report, and had no previous head injury with loss of consciousness or a history of seizures. Clinically stable patients with a DSM-IV diagnosis of schizophrenia were recruited from 5 sites across Ireland. Control participants were included only if they were aged 18 to 65 years and satisfied, based on clinical interview, the criteria of having no history of major mental health problems, intellectual disability, or a first-degree relative with a history of psychosis.

Clinical/Neuropsychological Assessment. Diagnosis was confirmed by trained psychiatrists using the SCID-I (First et al., 1995b). Additional diagnostic details and clinical sample characteristics ascertained at the time of interview included symptom severity as measured with SANS/SAPS (Andreasen, 1983, 1984) or PANSS (Kay et al., 1987), overall functioning using GAF (American Psychiatric Association, 1994), and medication dosage. Premorbid IQ was measured using the Wechsler Test of Adult Reading (Wechsler, 2001). Current IQ was measured using selected subtests (Vocabulary, Similarities, Block Design, and Matrix Reasoning) from the UK version of the WAIS-III (Wechsler, 2005). Speed of processing was assessed using the Trail Making Test—Part A and B (Reitan, 1958). Phonemic Verbal Fluency was assessed using the COWAT (Reitan, 1985) (number of words generated in 60 seconds that start on letter F, A and S). Attention/vigilance was measured using the 1-9 Continuous Performance Test (Gordon and Mettleman, 1987), or in a small subset using the three-letter condition of the CPT-IP (Cornblatt et al., 1988). Verbal working memory was measured using the Letter-Number Sequencing subtest from the Wechsler Memory Scale, Third Edition (WMS-III; Wechsler, 1997b) and spatial working memory using the Spatial Working Memory test from the CANTAB (Cambridge Cognition, 2003b) and a spatial n-back task (Callicott et al., 1998). Verbal learning and memory was assessed using the CVLT-II Short Form (Delis et al., 2000). Episodic memory was measured using the Logical Memory immediate and delayed tests from the WMS-III (Wechsler, 1997b). Visual learning and memory was assessed using the WMS-III Faces subtest (Wechsler, 1997b) and the CANTAB Paired Associates Learning subtest. Executive Functioning was assessed using the CANTAB Intra-Extra Dimensional Set Shifting subtest (Cambridge Cognition, 2003b) and the Sustained Attention to Response Task (Robertson et al., 1997). Finally, social cognition was measured using the Hinting Task (Versmissen et al., 2008) and Reading the Mind in the Eyes Test (Baron-Cohen et al., 2001).

MRI Acquisition. For the subsample recruited from Trinity College Dublin, T1-weighted structural scans (TE = 3.8 ms; TR = 8.4 ms; Flip angle = 8°; FOV = 23 cm; Spatial resolution = 0.9 x 0.9 x 0.9 mm; matrix = 256 x 256 mm; Slice acquisition direction = axial) and Diffusion Tensor Imaging (TE = 52 ms; TR = 12445 ms; Flip angle = 90°; FOV = 22.4 cm; Spatial resolution = 1.75 x 1.75 x 2.2 mm; matrix = 128 x 128 mm; Slice acquisition direction = axial; number of

diffusion-encoding gradient directions = 15; number of b0 volumes = 1; b-value = 800 s/mm²) were acquired on a 3 Tesla Philips Intera Achieva scanner at the Trinity College Institute for Neuroscience using a SENSE 8-channel head coil. For the subsample recruited from National University of Ireland Galway, T1-weighted structural scans (TE = 4.38 ms; TI = 600 ms; TR = 1140 ms; Flip angle = 15°; FOV = 23 cm; Spatial resolution = 0.45 x 0.45 x 0.9 mm; matrix = 512 x 512 mm; Slice acquisition direction = axial) were acquired on a 1.5 Tesla Siemens Magnetom Symphony scanner using a 4-channel head coil.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on the Affymetrix 6.0 or Illumina HumanCore Exome arrays.

Ethics/Consent. Assessments of all patient and control participants were conducted in accordance with the relevant ethics committee approval from each participating site. All patients and controls were of Irish ancestry (4 grandparents born in Ireland), and all provided written informed consent.

Acknowledgements. Recruitment and genotyping were supported by the Wellcome Trust and Science Foundation Ireland (SFI08/IN.1/B1916, A Corvin). Dr Donohoe is also supported by the European Research Council (ERC-2015-STG-677467). The investigators thank Prof. John Waddington, Prof. Ted Dinan, Prof. Eadbhard O'Callaghan, Prof. Kieran Murphy, and Dr. F. Anthony O'Neil for their role in the recruitment of the patient sample.

UMCU-SZ1 — University Medical Centre Utrecht, Schizophrenia study 1

(Hulshoff Pol et al., 2001; Terwisscha van Scheltinga et al., 2012; Terwisscha van Scheltinga et al., 2013a; Terwisscha van Scheltinga et al., 2013b)

Recruitment. Patients with schizophrenia or schizoaffective disorder were recruited from various outpatient and inpatient clinics; treatment setting was unrelated to age. Comparison control participants were matched with the patients for the socioeconomic status of their parents, which was expressed as the highest level of education completed by one parent. Participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014) and the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Schizophrenia Working Group dataset (van Erp et al., 2016).

Inclusion/Exclusion Criteria. All participants were between 16 and 70 years of age. Participants with a major medical or neurological illness, including past head trauma, hypertension, cardiac disease, diabetes mellitus, cerebrovascular disease, epilepsy, migraine, endocrine disorders, drug or alcohol dependence, or an IQ below 80 were excluded. Control participants had no history of psychiatric illness except for three participants who had a history of anxiety disorder, obsessive-compulsive disorder and adjustment disorder, respectively. None of the control participants had first-degree family members with psychotic illness.

Clinical/Neuropsychological Assessment. Clinical diagnosis was determined using the Comprehensive Assessment of Symptoms and History (CASH; Andreasen et al., 1992)) and the Schedule for Affective Disorders and Schizophrenia (Endicott and Spitzer, 1978) and was assessed by two independent raters. Diagnostic consensus was achieved in the presence of a psychiatrist. The PANSS (Kay et al., 1987) and GAF (American Psychiatric Association, 1994) were used to measure symptom levels and overall functioning, respectively. Premorbid IQ was estimated using the Dutch version of the NART (Schmand et al., 1991). Current IQ was estimated with the Dutch version of the WAIS-III (Stinissen et al., 1970; Wechsler, 2000). Four out of 11 subtests of WAIS-III were used: Block Design, Comprehension, Vocabulary and Picture arrangement. Raw scores were multiplied by 11/4 to obtain an estimate of total IQ. Verbal fluency was assessed using the Multilingual Aphasia Examination (Benton and Hamsher, 1978) including a Phonemic Verbal Fluency task (number of words generated in 60 seconds that start on letter N, A and P) and a Semantic Verbal Fluency task (category: animals). Sustained visual attention and vigilance were assessed using the Continuous Performance Task-HQ (CPT A-X; Nuechterlein and Dawson, 1984)), with outcome parameters of reaction time and accuracy. Verbal learning

and memory was assessed with the CVLT (Delis et al., 1987). Episodic memory was assessed using the Logical Memory subtests from the Dutch WMS-R (Wechsler, 1987). Visual learning memory was assessed using the WMS-R Visual Reproduction subtests. Executive functioning was assessed using the Stroop test (Stroop, 1935).

MRI Acquisition. T1-weighted structural scans (TE = 4.6 ms; TR = 30 ms; Flip angle = 30°; FOV = 25.6 cm; Spatial resolution = 1.0 x 1.0 x 1.2 mm; matrix = 256 x 256 mm; Slice acquisition direction = coronal) were acquired on a 1.5 Tesla Philips NT Intera scanner using a 6-element SENSE receiver head coil.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on the Illumina HumanHap550 or Illumina Infinium OmniExpressExome-8 arrays. Genotype data are available from the NIMH Repository and Genomics Resource (<http://www.nimhgenetics.org>).

Ethics/Consent. The study was approved by the Medical Ethical Committee of the University Medical Center Utrecht.

Acknowledgements. The study was supported by grants 7F99.(2).37 (H Hulshoff Pol) and 14F06(2)-34 (SC Bakker) from the HersenStichting Nederland (Dutch Brain Foundation), The Hague, the Netherlands, by a Veni grant from Zorg Onderzoek Nederland, Medische Wetenschappen to SC Bakker (the Dutch organization for health research and development, project number: 91686137), and by Top Institute Pharma (project T5-203). Genome-wide SNP genotyping was funded by NIMH grant R01MH078075 (RA Ophoff).

UMCU-SZ2 — University Medical Centre Utrecht, Schizophrenia study 2

(Boos et al., 2013; Korver et al., 2012; Rais et al., 2012; Terwisscha van Scheltinga et al., 2013a)

Recruitment. Data pertain to baseline measures of a longitudinal study (GROUP) in the Netherlands and Belgium. In selected representative geographical areas, patients were identified through clinicians working in regional psychotic disorder services whose caseloads were screened for inclusion criteria. Subsequently, a group of patients presenting consecutively at these services as either outpatients or in-patients were recruited for the study. Controls were selected through a system of random mailings to addresses in the catchment areas of the patients. Participants form part of the Psychiatric Genomics Consortium Schizophrenia Working Group dataset (PGC Schizophrenia Working Group, 2014).

Inclusion/Exclusion Criteria. Inclusion criteria for patients, siblings, and controls were: (1) age range of 16–50 years and (2) good command of the Dutch language. Patients had to meet Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR) criteria for a non-affective psychotic disorder (American Psychiatric Association, 2000). Exclusion criteria for healthy controls were a history of psychotic disorder or a first-degree family member with a history of psychotic disorder.

Clinical/Neuropsychological Assessment. Diagnosis was assessed with the Comprehensive Assessment of Symptoms and History interview (CASH; Andreasen et al., 1992) or the Schedules for Clinical Assessment for Neuropsychiatry (SCAN) version 2.1 (Wing et al., 1990). The PANSS (Kay et al., 1987) and GAF (American Psychiatric Association, 1994) were used to measure symptom levels and overall functioning, respectively. Additionally, the Structured Interview for Schizotypy-Revised (SIS-R; Vollema and Ormel, 2000) and the CDSS (Addington et al., 1992) were administered. The cognitive assessment took between 90 and 120 min. The following subtests of the WAIS-III (Wechsler, 1997a) were assessed: Digit Symbol Coding as a measure of processing speed; Arithmetic as a measure of working memory; Information as a measure of acquired knowledge; and Block Design as a measure of reasoning and problem solving. Together, these 3 subtests were used to estimate Current IQ. Sustained visual attention and vigilance were assessed using the Continuous Performance Task-H-Q, a variation of the CPT A-X by Nuechterlein and Dawson, 1984), with outcome parameters of reaction time and accuracy. Verbal learning was assessed using the Auditory Verbal Learning Test (Brand and Jolles, 1985), with

outcome parameters of immediate recall (15-word list, three learning trials) and retention rate after 20 min. Set shifting ability was assessed using the Response Shifting Task, a modified version of the Competing Programs Task (Bilder et al., 1992; Nolan et al., 2004), with outcome parameters of reaction time and accuracy. The Degraded Facial Affect Recognition Task (van 't Wout et al., 2004) was used to assess recognition of neutral, happy, fearful and angry emotions. The Benton Face Recognition Task (Benton et al., 1983) was used to assess visuospatial discrimination of unfamiliar faces. The Hinting Task (Versmissen et al., 2008) was used to assess theory of mind.

MRI Acquisition. T1-weighted structural scans (TE = 4.6 ms; TR = 30 ms; Flip angle = 30°; FOV = 25.6 cm/70%; Spatial resolution = 1.0 x 1.0 x 1.2 mm; matrix = 256 x 256 mm; Slice acquisition direction = coronal) and Diffusion Tensor Imaging (TE = 88 ms, TR = 9822 ms; Flip angle = 90°; FOV = 24 cm; Spatial resolution = 2.0 x 2.0 x 2.0 mm; matrix = 128 x 96 mm; Slice acquisition direction = axial; number of diffusion-encoding gradient directions = 32; number of b0 volumes = 8; b-value = 1000 s/mm²) were acquired on a 1.5 Tesla Philips Achieva scanner using a 6-element SENSE receiver head coil.

Genotyping. DNA from a proportion of participants was obtained from whole blood and genotyped on the Affymetrix 6.0, Illumina HumanHap550, or Illumina Infinium OmniExpressExome-8 arrays. Genotype data are available from the NIMH Repository and Genomics Resource (<http://www.nimhgenetics.org>).

Ethics/Consent. The study protocol was approved centrally by the Ethical Review Board of the University Medical Center Utrecht and subsequently by local review boards of each participating institute. All participants gave written informed consent in accordance with the committee's guidelines.

Acknowledgements. The study was funded through the Geestkracht programme of the Dutch Health Research Council (ZON-MW, grant number 10-000-1001), and matching funds from participating industry (Lundbeck, AstraZeneca, Eli Lilly, Janssen Cilag), universities and mental health care organizations in the Netherlands. Genome-wide SNP genotyping was funded by NIMH grant R01MH078075 (RA Ophoff).

ZHH — Zucker Hillside Hospital

(Lencz et al., 2014)

Recruitment. Healthy participants from the New York metropolitan area were recruited through advertisements, word of mouth, referrals, and study registries. Participants form part of the Cognitive Genomics Consortium wave 1 (COGENT1) dataset (Lencz et al., 2014).

Inclusion/Exclusion Criteria. Participants had no history of a current or past DSM-IV axis I major mood or psychotic disorder as assessed by structured diagnostic interview (First et al., 1995a). Other exclusion criteria included: intellectual or learning disability; significant medical illness that could affect brain structure; or lifetime exposure to psychiatric medication.

Clinical/Neuropsychological Assessment. Premorbid IQ was estimated using the WRAT-3 (Wilkinson, 1993). Further cognitive testing was performed using the MATRICS Consensus Cognitive Battery (MCCB) (Kern et al., 2008; Nuechterlein et al., 2008). Speed of processing was assessed using the BACS Symbol Coding subtest (Keefe et al., 2008), the WAIS-R Digit Symbol subtest (Wechsler, 1981), Trail Making Test—Part A and B (Reitan, 1958), Semantic Verbal Fluency (MCCB Category Fluency; category: animals) (Kern et al., 2008; Nuechterlein et al., 2008), and COWAT Phonemic Verbal Fluency (letters F, A, S) (Benton and Hamsher, 1978; Reitan, 1985); attention using the CPT-IP (Cornblatt et al., 1988); working memory using the WMS-III Spatial Span subtest (Wechsler, 1997b), University of Maryland Letter-Number Span (Gold et al., 1997), and the WAIS-R Digit Span subtest (Wechsler, 1981); verbal learning using the HVLIT-R (Brandt and Benedict, 1997); visual learning using the BVMT-R (Benedict, 1997); reasoning and problem solving using the WCST (Heaton et al., 1993) and NAB Mazes (Stern and White, 2003); and social cognition using the MSCEIT (Mayer et al., 2003).

MRI Acquisition. Not applicable.

Genotyping. DNA obtained from whole blood was genotyped on the Illumina OmniExpress array.

Ethics/Consent. This study was approved by the Institutional Review Board of the North Shore – Long Island Jewish Health System. Written informed consent was obtained from all participants.

Acknowledgements. The study was supported by the Donald and Barbara Zucker Foundation, internal funding from the North Shore-Long Island Jewish Health System, and grants from the Stanley Foundation (EJ Bromet), the National Alliance for Research on Schizophrenia and Depression (AK Malhotra), and the NIMH (K23MH001760, R01MH079800, and P50MH080173 to AK Malhotra, R01MH0084098 and RC2MH089964 to T Lencz, R01MH044801 to EJ Bromet, R01MH060374 to RM Bilder, and Center Grants P30MH074543 to JM Kane and M01RR018535 to KJ Tracey).

Conflicts of interest. Dr. Lencz has received consulting fees or honoraria from Merck, Eli Lilly & Company, Cowen & Co., Guidepoint Global, Golden Helix, InforMed Insights, and PGxHealth (a division of Clinical Data). Dr Malhotra has received consulting fees or honoraria from Eli Lilly & Company, Janssen Pharmaceutica, Merck, Bristol-Meyers Squibb, Pfizer, PGxHealth (a division of Clinical Data), Roche Diagnostics, and Vanda Pharmaceuticals, and has received research support from Eli Lilly & Company.

References

- Addington, D., Addington, J., Maticka-Tyndale, E., Joyce, J., 1992. Reliability and validity of a depression rating scale for schizophrenics. *Schizophr Res* **6** (3), 201-208. PMID: 1571313.
- American Psychiatric Association, 1994. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV). American Psychiatric Press Inc., Washington, DC.
- American Psychiatric Association, 2000. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR). American Psychiatric Press Inc., Washington, DC.
- Andreasen, N.C., 1983. Scale for the Assessment of Negative Symptoms (SANS). University of Iowa, Iowa City.
- Andreasen, N.C., 1984. Scale for the Assessment of Positive Symptoms (SAPS). University of Iowa, Iowa City.
- Andreasen, N.C., Flaum, M., Arndt, S., 1992. The Comprehensive Assessment of Symptoms and History (CASH). An instrument for assessing diagnosis and psychopathology. *Arch Gen Psych* **49** (8), 615-623. PMID: 1637251.
- Annett, M., 1970. A classification of hand preference by association analysis. *Br J Psychol* **61** (3), 303-321. PMID: 5457503.
- Army Individual Test Battery, 1944. Army Individual Test Battery: Manual of directions and scoring. War Department, Adjutant General's Office, Washington, DC.
- Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., Plumb, I., 2001. The "Reading the Mind in the Eyes" Test revised version: A study with normal adults, and adults with Asperger syndrome or high-functioning autism. *J Child Psychol Psychiatry* **42** (2), 241-251. PMID: 11280420.
- Benedict, R.H.B., 1997. Brief Visuospatial Memory Test-Revised. Psychological Assessment Resources, Inc., Odessa, FL.
- Benton, A.L., Hamsher, K.d., 1978. Multilingual Aphasia Examination Manual (revised). University of Iowa, Iowa City, IA.
- Benton, A.L., Hamsher, K.d., 1989. Multilingual Aphasia Examination. AJA Associates, Iowa City, IA.
- Benton, A.L., Hamsher, K.d., Varney, N.R., Spreen, O., 1983. Contributions to Neuropsychological Assessment: A Clinical Manual. Oxford University Press, New York, NY.
- Benton, A.L., Sivan, A.B., Hamsher, K.d., Varney, N.R., Spreen, O., 1994. Contributions to Neuropsychological Assessment: A Clinical Manual. Oxford University Press, Oxford, UK.
- Bhojraj, T.S., Francis, A.N., Montrose, D.M., Keshavan, M.S., 2011. Grey matter and cognitive deficits in young relatives of schizophrenia patients. *Neuroimage* **54 Suppl 1**, S287-292. PMID: 20362681.
- Bhojraj, T.S., Francis, A.N., Rajarethinam, R., Eack, S., Kulkarni, S., Prasad, K.M., Montrose, D.M., Dworakowski, D., Diwadkar, V., Keshavan, M.S., 2009. Verbal fluency deficits and altered lateralization of language brain areas in individuals genetically predisposed to schizophrenia. *Schizophr Res* **115** (2-3), 202-208. PMID: 19840895.
- Bilder, R.M., Turkel, E., Lipschutz-Broch, L., Lieberman, J.A., 1992. Antipsychotic medication effects on neuropsychological functions. *Psychopharmacol Bull* **28** (4), 353-366. PMID: 1363583.
- Blair, J.R., Spreen, O., 1989. Predicting premorbid IQ: A revision of the National Adult Reading Test. *The Clinical Neuropsychologist* **3** (2), 129-136. PMID: N/A.
- Boos, H.B., Mandl, R.C., van Haren, N.E., Cahn, W., van Baal, G.C., Kahn, R.S., Hulshoff Pol, H.E., 2013. Tract-based diffusion tensor imaging in patients with schizophrenia and their non-psychotic siblings. *Eur Neuropsychopharmacol* **23** (4), 295-304. PMID: 22841128.
- Brand, N., Jolles, J., 1985. Learning and retrieval rate of words presented auditorily and visually. *J Gen Psychol* **112** (2), 201-210. PMID: 4056765.
- Brandt, J., 1991. The Hopkins Verbal Learning Test: Development of a new memory test with six equivalent forms. *Clin Neuropsychol* **5** (2), 125-142. PMID: N/A.
- Brandt, J., Benedict, R.H.B., 1997. Hopkins Verbal Learning Test-Revised. Psychological Assessment Resources, Inc., Odessa, FL.
- Buka, S.L., Seidman, L.J., Tsuang, M.T., Goldstein, J.M., 2013. The New England Family Study High-risk Project: neurological impairments among offspring of parents with schizophrenia and other psychoses. *Am J Med Genet B Neuropsychiatr Genet* **162B** (7), 653-660. PMID: 24132897.
- Callicott, J.H., Ramsey, N.F., Tallent, K., Bertolino, A., Knable, M.B., Coppola, R., Goldberg, T., van Gelderen, P., Mattay, V.S., Frank, J.A., Moonen, C.T., Weinberger, D.R., 1998. Functional magnetic resonance imaging brain mapping in psychiatry: Methodological issues illustrated in a study of working memory in schizophrenia. *Neuropsychopharmacology* **18** (3), 186-196. PMID: 9471116.
- Cambridge Cognition, 2003a. Cambridge Neuropsychological Test Automated Battery (CANTAB). Cambridge Cognition Ltd, Cambridge, UK.
- Cambridge Cognition, 2003b. Cambridge Neuropsychological Test Automated Battery, expedio version (CANTABexpedio). Cambridge Cognition Ltd, Cambridge, UK.
- Clemm von Hohenberg, C., Pasternak, O., Kubicki, M., Ballinger, T., Vu, M.A., Swisher, T., Green, K., Giwerc, M., Dahlben, B., Goldstein, J.M., Woo, T.U., Petryshen, T.L., Meshulam-Gately, R.I., Woodberry, K.A., Thermenos, H.W., Muler, C., McCarley, R.W., Seidman, L.J., Shenton, M.E., 2014. White matter microstructure in individuals at clinical high risk of psychosis: A whole-brain diffusion tensor imaging study. *Schizophr Bull* **40** (4), 895-903. PMID: 23737549.

- Coleman, A.R., Moberg, P.J., Ragland, J.D., Gur, R.C., 1997. Comparison of the Halstead-Reitan and Infrared Light Beam Finger Tappers. *Assessment* **4** (3), 277-286. PMID: 26613776.
- Cornblatt, B.A., Risch, N.J., Faris, G., Friedman, D., Erlenmeyer-Kimling, L., 1988. The Continuous Performance Test, Identical Pairs version (CPT-IP): I. New findings about sustained attention in normal families. *Psychiatry Res* **26** (2), 223-238. PMID: 3237915.
- del Re, E.C., Bergen, S.E., Mesholam-Gately, R.I., Niznikiewicz, M.A., Goldstein, J.M., Woo, T.U., Shenton, M.E., Seidman, L.J., McCarley, R.W., Petryshen, T.L., 2014. Analysis of schizophrenia-related genes and electrophysiological measures reveals ZNF804A association with amplitude of P300b elicited by novel sounds. *Transl Psychiatry* **4**, e346. PMID: 24424392.
- Delis, D.C., Kaplan, E., Kramer, J.H., 2001. Delis-Kaplan Executive Function System. The Psychological Corporation, San Antonio, TX.
- Delis, D.C., Kramer, J.H., Kaplan, E., Holdnack, J., 2004. Reliability and validity of the Delis-Kaplan Executive Function System: An update. *J Int Neuropsychol Soc* **10** (2), 301-303. PMID: 15012851.
- Delis, D.C., Kramer, J.H., Kaplan, E., Ober, B.A., 1987. California Verbal Learning Test-Adult version. The Psychological Corporation, New York, NY.
- Delis, D.C., Kramer, J.H., Kaplan, E., Ober, B.A., 2000. California Verbal Learning Test: Second Edition. Psychological Corporation, San Antonio, TX.
- Denckla, M.B., Rudel, R., 1974. Rapid "automatized" naming of pictured objects, colors, letters and numbers by normal children. *Cortex* **10** (2), 186-202. PMID: 4844470.
- Di Forti, M., Morgan, C., Dazzan, P., Pariante, C., Mondelli, V., Marques, T.R., Handley, R., Luzi, S., Russo, M., Paparelli, A., Butt, A., Stilo, S.A., Wiffen, B., Powell, J., Murray, R.M., 2009. High-potency cannabis and the risk of psychosis. *Br J Psychiatry* **195** (6), 488-491. PMID: 19949195.
- Donohoe, G., Walters, J., Morris, D.W., Quinn, E.M., Judge, R., Norton, N., Giegling, I., Hartmann, A.M., Moller, H.J., Muglia, P., Williams, H., Moskvina, V., Peel, R., O'Donoghue, T., Owen, M.J., O'Donovan, M.C., Gill, M., Rujescu, D., Corvin, A., 2009. Influence of NOS1 on verbal intelligence and working memory in both patients with schizophrenia and healthy control subjects. *Arch Gen Psychiatry* **66** (10), 1045-1054. PMID: 19805695.
- Endicott, J., Spitzer, R.L., 1978. A diagnostic interview: The Schedule for Affective Disorders and Schizophrenia. *Arch Gen Psychiatry* **35** (7), 837-844. PMID: 678037.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B., 1994. Structured Clinical Interview for DSM-IV Axis I Disorders—Administration Booklet. American Psychiatric Press Inc., Washington, DC.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B., 1995a. Structured Clinical Interview for DSM-IV Axis I Disorders—Non-Patient Edition (SCID-I/NP). Biometrics Research Department, New York State Psychiatric Institute, New York, NY.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B., 1995b. Structured Clinical Interview for DSM-IV Axis I Disorders—Patient Edition (SCID-I/P, Version 2.0). Biometrics Research Department, New York State Psychiatric Institute, New York, NY.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B.W., 2002a. Structured Clinical Interview for DSM-IV-TR Axis I Disorders—Research Version, Non-Patient Edition. Biometrics Research Department, New York State Psychiatric Institute, New York, NY.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B.W., 2002b. Structured Clinical Interview for DSM-IV-TR Axis I Disorders—Research Version, Patient Edition. Biometrics Research Department, New York State Psychiatric Institute, New York, NY.
- Folstein, M.F., Robins, L.N., Helzer, J.E., 1983. The Mini-Mental State Examination. *Arch Gen Psychiatry* **40** (7), 812. PMID: 6860082.
- Francis, A.N., Seidman, L.J., Jabbar, G.A., Mesholam-Gately, R., Thermenos, H.W., Juelich, R., Proal, A.C., Shenton, M., Kubicki, M., Mathew, I., Keshavan, M., Delisi, L.E., 2012. Alterations in brain structures underlying language function in young adults at high familial risk for schizophrenia. *Schizophr Res* **141** (1), 65-71. PMID: 22892286.
- Glahn, D.C., Gur, R.C., Ragland, J.D., Censits, D.M., Gur, R.E., 1997. Reliability, performance characteristics, construct validity, and an initial clinical application of a visual object learning test (VOLT). *Neuropsychology* **11** (4), 602-612. PMID: 9345704.
- Gold, J.M., Carpenter, C., Randolph, C., Goldberg, T.E., Weinberger, D.R., 1997. Auditory working memory and Wisconsin Card Sorting Test performance in schizophrenia. *Arch Gen Psychiatry* **54** (2), 159-165. PMID: 9040284.
- Goldstein, J.M., Buka, S.L., Seidman, L.J., Tsuang, M.T., 2010. Specificity of familial transmission of schizophrenia psychosis spectrum and affective psychoses in the New England Family Study's high-risk design. *Arch Gen Psychiatry* **67** (5), 458-467. PMID: 20439827.
- Goldstein, J.M., Cherkertzian, S., Seidman, L.J., Donatelli, J.A., Remington, A.G., Tsuang, M.T., Hornig, M., Buka, S.L., 2014. Prenatal maternal immune disruption and sex-dependent risk for psychoses. *Psychol Med* **44** (15), 3249-3261. PMID: 25065485.
- Gollub, R.L., Shoemaker, J.M., King, M.D., White, T., Ehrlich, S., Sponheim, S.R., Clark, V.P., Turner, J.A., Mueller, B.A., Magnotta, V., O'Leary, D., Ho, B.C., Brauns, S., Manoach, D.S., Seidman, L., Bustillo, J.R., Lauriello, J., Bockholt, J., Lim, K.O., Rosen, B.R., Schulz, S.C., Calhoun, V.D., Andreasen, N.C., 2013. The MCIC collection: A

- shared repository of multi-modal, multi-site brain image data from a clinical investigation of schizophrenia. *Neuroinformatics* **11** (3), 367-388. PMID: 23760817.
- Gordon, M., Mettleman, B.B., 1987. Technical Guide to the Gordon Diagnostic System (GDS). Gordon Systems, DeWitt, NY.
- Gur, R.C., Richard, J., Hughett, P., Calkins, M.E., Macy, L., Bilker, W.B., Brensinger, C., Gur, R.E., 2010. A cognitive neuroscience-based computerized battery for efficient measurement of individual differences: Standardization and initial construct validation. *J Neurosci Methods* **187** (2), 254-262. PMID: 19945485.
- Gur, R.C., Sara, R., Hagendoorn, M., Marom, O., Hughett, P., Macy, L., Turner, T., Bajcsy, R., Posner, A., Gur, R.E., 2002. A method for obtaining 3-dimensional facial expressions and its standardization for use in neurocognitive studies. *J Neurosci Methods* **115** (2), 137-143. PMID: 11992665.
- Guy, W., 2000. Clinical Global Impressions (CGI) Scale, Modified, in: Rush, J.A. (Ed.), Task Force for the Handbook of Psychiatric Measures. Handbook of Psychiatric Measures, 1st ed. American Psychiatric Association, Washington, DC.
- Hamilton, M., 1960. A rating scale for depression. *J Neurol Neurosurg Psychiatry* **23**, 56-62. PMID: 14399272.
- Härting, C., Markowitsch, H.J., Neufeld, H., Calabrese, P., Deisinger, K., Kessler, J., 2000. Wechsler Gedächtnistest–Revidierte Fassung [WMS-R, German version]. Huber, Bern, Switzerland.
- Heaton, R.K., 2003. Wisconsin Card Sorting Test: Computer Version 4 – Research edition. Psychological Assessment Resources, Inc., Odessa, FL.
- Heaton, R.K., Chelune, G.J., Taley, J.L., Kay, G.G., Curtiss, G., 1993. Wisconsin Card Sorting Test Manual: Revised and Expanded. Psychological Assessment Resources, Odessa, FL.
- Helmstaedter, C., Lendt, M., Lux, S., 2001. Verbaler Lern- und Merkfähigkeitstest (VLMT) [Verbal Learning and Memory Test]. Beltz, Göttingen, Germany.
- Ho, N.F., Iglesias, J.E., Sum, M.Y., Kuswanto, C.N., Sitoh, Y.Y., De Souza, J., Hong, Z., Fischl, B., Roffman, J.L., Zhou, J., Sim, K., Holt, D.J., 2016. Progression from selective to general involvement of hippocampal subfields in schizophrenia. *Mol Psychiatry* **22** (1), 142-152. PMID: 26903271.
- Hollingshead, A.A., 1975. Four-factor index of social status, Yale University, New Haven, CT.
- Holt, D.J., Coombs, G., Zeidan, M.A., Goff, D.C., Milad, M.R., 2012. Failure of neural responses to safety cues in schizophrenia. *Arch Gen Psychiatry* **69** (9), 893-903. PMID: 22945619.
- Hulshoff Pol, H.E., Schnack, H.G., Mandl, R.C., van Haren, N.E., Koning, H., Collins, D.L., Evans, A.C., Kahn, R.S., 2001. Focal gray matter density changes in schizophrenia. *Arch Gen Psychiatry* **58** (12), 1118-1125. PMID: 11735840.
- Kay, S.R., Fiszbein, A., Opler, L.A., 1987. The Positive And Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophr Bull* **13** (2), 261-276. PMID: 3616518.
- Keefe, R.S., Bilder, R.M., Harvey, P.D., Davis, S.M., Palmer, B.W., Gold, J.M., Meltzer, H.Y., Green, M.F., Miller, D.D., Canive, J.M., Adler, L.W., Manschreck, T.C., Swartz, M., Rosenheck, R., Perkins, D.O., Walker, T.M., Stroup, T.S., McEvoy, J.P., Lieberman, J.A., 2006. Baseline neurocognitive deficits in the CATIE schizophrenia trial. *Neuropsychopharmacology* **31** (9), 2033-2046. PMID: 16641947.
- Keefe, R.S., Goldberg, T.E., Harvey, P.D., Gold, J.M., Poe, M.P., Coughenour, L., 2004. The Brief Assessment of Cognition in Schizophrenia: Reliability, sensitivity, and comparison with a standard neurocognitive battery. *Schizophr Res* **68** (2-3), 283-297. PMID: 15099610.
- Keefe, R.S., Harvey, P.D., Goldberg, T.E., Gold, J.M., Walker, T.M., Kennel, C., Hawkins, K., 2008. Norms and standardization of the Brief Assessment of Cognition in Schizophrenia (BACS). *Schizophr Res* **102** (1-3), 108-115. PMID: 18495435.
- Kendler, K.S., Lieberman, J.A., Walsh, D., 1989. The Structured Interview for Schizotypy (SIS): A preliminary report. *Schizophr Bull* **15** (4), 559-571. PMID: 2623438.
- Kern, R.S., Nuechterlein, K.H., Green, M.F., Baade, L.E., Fenton, W.S., Gold, J.M., Keefe, R.S., Meshulam-Gately, R., Mintz, J., Seidman, L.J., Stover, E., Marder, S.R., 2008. The MATRICS Consensus Cognitive Battery, part 2: Co-norming and standardization. *Am J Psychiatry* **165** (2), 214-220. PMID: 18172018.
- Kerr, S.L., Neale, J.M., 1993. Emotion perception in schizophrenia: Specific deficit or further evidence of generalized poor performance? *J Abnorm Psychol* **102** (2), 312-318. PMID: 8315144.
- Kikinis, Z., Fallon, J.H., Niznikiewicz, M., Nestor, P., Davidson, C., Bobrow, L., Pelavin, P.E., Fischl, B., Yendiki, A., McCarley, R.W., Kikinis, R., Kubicki, M., Shenton, M.E., 2010. Gray matter volume reduction in rostral middle frontal gyrus in patients with chronic schizophrenia. *Schizophr Res* **123** (2-3), 153-159. PMID: 20822884.
- Kikinis, Z., Fitzsimmons, J., Dunn, C., Vu, M.A., Makris, N., Bouix, S., Goldstein, J.M., Meshulam-Gately, R.I., Petryshen, T., Del Re, E.C., Wojcik, J., Seidman, L.J., Kubicki, M., 2015. Anterior commissural white matter fiber abnormalities in first-episode psychosis: A tractography study. *Schizophr Res* **162** (1-3), 29-34. PMID: 25667192.
- Korver, N., Quee, P.J., Boos, H.B., Simons, C.J., de Haan, L., GROUP investigators, 2012. Genetic Risk and Outcome of Psychosis (GROUP), a multi-site longitudinal cohort study focused on gene-environment interaction: Objectives, sample characteristics, recruitment and assessment methods. *Int J Methods Psychiatr Res* **21** (3), 205-221. PMID: 22419500.
- Kuswanto, C.N., Sum, M.Y., Qiu, A., Sitoh, Y.Y., Liu, J., Sim, K., 2015. The impact of genome wide supported microRNA-137 (MIR137) risk variants on frontal and striatal white matter integrity, neurocognitive functioning, and

- negative symptoms in schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* **168B** (5), 317-326. PMID: 25921703.
- Lafayette Instrument Company -- Trites, R., 1989. Grooved Pegboard Instruction Manual, Model 32025. Lafayette Instrument Company, Lafayette, IN.
- Lam, M., Collinson, S.L., Eng, G.K., Rapisarda, A., Kraus, M., Lee, J., Chong, S.A., Keefe, R.S., 2014. Refining the latent structure of neuropsychological performance in schizophrenia. *Psychol Med* **44** (16), 3557-3570. PMID: 25066336.
- Lencz, T., Knowles, E., Davies, G., Guha, S., Liewald, D.C., Starr, J.M., Djurovic, S., Melle, I., Sundet, K., Christoforou, A., Reinvang, I., Mukherjee, S., DeRosse, P., Lundervold, A., Steen, V.M., John, M., Espeseth, T., Raikkonen, K., Widen, E., Palotie, A., Eriksson, J.G., Giegling, I., Konte, B., Ikeda, M., Roussos, P., Giakoumaki, S., Burdick, K.E., Payton, A., Ollier, W., Horan, M., Donohoe, G., Morris, D., Corvin, A., Gill, M., Pendleton, N., Iwata, N., Darvasi, A., Bitsios, P., Rujescu, D., Lahti, J., Hellard, S.L., Keller, M.C., Andreassen, O.A., Deary, I.J., Glahn, D.C., Malhotra, A.K., 2014. Molecular genetic evidence for overlap between general cognitive ability and risk for schizophrenia: A report from the Cognitive Genomics consortium (COGENT). *Mol Psychiatry* **19** (2), 168-174. PMID: 24342994.
- Lieberman, J.A., 2007. Effectiveness of antipsychotic drugs in patients with chronic schizophrenia: Efficacy, safety and cost outcomes of CATIE and other trials. *J Clin Psychiatry* **68** (2), e04. PMID: 17335312.
- Linnman, C., Coombs, G., 3rd, Goff, D.C., Holt, D.J., 2013. Lack of insula reactivity to aversive stimuli in schizophrenia. *Schizophr Res* **143** (1), 150-157. PMID: 23201307.
- Lyons-Warren, A., Lillie, R., Hershey, T., 2004. Short- and long-term spatial delayed response performance across the lifespan. *Dev Neuropsychol* **26** (3), 661-678. PMID: 15525563.
- Mallett, R., Leff, J., Bhugra, D., Pang, D., Zhao, J.H., 2002. Social environment, ethnicity and schizophrenia. A case-control study. *Soc Psychiatry Psychiatr Epidemiol* **37** (7), 329-335. PMID: 12111025.
- Maxwell, M.E., 1996. FIGS: Clinical Neurogenetics Branch, Intramural Research Program. NIMH, Bethesda, MD.
- Mayer, J.D., Salovey, P., Caruso, D.R., Sitarenios, G., 2003. Measuring emotional intelligence with the MSCEIT V2.0. *Emotion* **3**, 97-105. PMID: 12899321.
- McClay, J.L., Adkins, D.E., Aberg, K., Bukszar, J., Khachane, A.N., Keefe, R.S., Perkins, D.O., McEvoy, J.P., Stroup, T.S., Vann, R.E., Beardsley, P.M., Lieberman, J.A., Sullivan, P.F., van den Oord, E.J., 2011. Genome-wide pharmacogenomic study of neurocognition as an indicator of antipsychotic treatment response in schizophrenia. *Neuropsychopharmacology* **36** (3), 616-626. PMID: 21107309.
- McGuffin, P., Farmer, A., Harvey, I., 1991. A polydiagnostic application of operational criteria in studies of psychotic illness: Development and reliability of the OPCRIT system. *Arch Gen Psychiatry* **48**, 764-770. PMID: 1883262.
- Miller, E., 1990. California Computerized Assessment Battery (CalCAP). Norland Software.
- Miller, M.D., Paradis, C.F., Houck, P.R., Mazumdar, S., Stack, J.A., Rifai, A.H., Mulsant, B., Reynolds, C.F., 1992. Rating chronic medical illness burden in geropsychiatric practice and research: Application of the Cumulative Illness Rating Scale. *Psychiatry Res* **41** (3), 237-248. PMID: 1594710.
- Miller, T.J., McGlashan, T.H., Rosen, J.L., Cadenhead, K., Cannon, T., Ventura, J., McFarlane, W., Perkins, D.O., Pearson, G.D., Woods, S.W., 2003. Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, interrater reliability, and training to reliability. *Schizophr Bull* **29** (4), 703-715. PMID: 14989408.
- Miller, T.J., McGlashan, T.H., Woods, S.W., Stein, K., Driesen, N., Corcoran, C.M., Hoffman, R., Davidson, L., 1999. Symptom assessment in schizophrenic prodromal states. *Psychiatr Q* **70** (4), 273-287. PMID: 10587984.
- Moore, T.M., Reise, S.P., Gur, R.E., Hakonarson, H., Gur, R.C., 2015. Psychometric properties of the Penn Computerized Neurocognitive Battery. *Neuropsychology* **29** (2), 235-246. PMID: 25180981.
- Nazeri, A., Chakravarty, M.M., Felsky, D., Lobaugh, N.J., Rajji, T.K., Mulsant, B.H., Voineskos, A.N., 2013. Alterations of superficial white matter in schizophrenia and relationship to cognitive performance. *Neuropsychopharmacology* **38** (10), 1954-1962. PMID: 23591167.
- Nelson, H.E., 1982. The National Adult Reading Test (NART): Test Manual. NFER-Nelson, Windsor, UK.
- Nelson, H.E., Willison, J., 1991. The National Adult Reading Test (NART). NFER-Nelson, Windsor, UK.
- Nestor, P.G., Shenton, M.E., McCarley, R.W., Haimson, J., Smith, R.S., O'Donnell, B., Kimble, M., Kikinis, R., Jolesz, F.A., 1993. Neuropsychological correlates of MRI temporal lobe abnormalities in schizophrenia. *Am J Psychiatry* **150** (12), 1849-1855. PMID: 8238641.
- Niswander, K.R., Gordon, M., 1972. The Collaborative Perinatal Study of the National Institute of Neurological Diseases and Stroke: The women and their pregnancies. U.S. Department of Health, Education, and Welfare, Washington, DC.
- Nolan, K.A., Bilder, R.M., Lachman, H.M., Volavka, J., 2004. Catechol O-methyltransferase Val158Met polymorphism in schizophrenia: Differential effects of Val and Met alleles on cognitive stability and flexibility. *Am J Psychiatry* **161** (2), 359-361. PMID: 14754787.
- Nuechterlein, K.H., Asarnow, R., 2004. 3-7 Continuous Performance Test. University of California Los Angeles, Los Angeles, CA.
- Nuechterlein, K.H., Dawson, M.E., 1984. Information processing and attentional functioning in the developmental course of schizophrenic disorders. *Schizophr Bull* **10** (2), 160-203. PMID: 6729409.

- Nuechterlein, K.H., Green, M.F., Kern, R.S., Baade, L.E., Barch, D.M., Cohen, J.D., Essock, S., Fenton, W.S., Frese, F.J., 3rd, Gold, J.M., Goldberg, T., Heaton, R.K., Keefe, R.S., Kraemer, H., Mesholam-Gately, R., Seidman, L.J., Stover, E., Weinberger, D.R., Young, A.S., Zalcman, S., Marder, S.R., 2008. The MATRICS Consensus Cognitive Battery, part 1: Test selection, reliability, and validity. *Am J Psychiatry* **165** (2), 203-213. PMID: 18172019.
- Nurnberger, J.I., Jr., Blehar, M.C., Kaufmann, C.A., York-Cooler, C., Simpson, S.G., Harkavy-Friedman, J., Severe, J.B., Malaspina, D., Reich, T., 1994. Diagnostic interview for genetic studies. Rationale, unique features, and training. NIMH Genetics Initiative. *Arch Gen Psychiatry* **51** (11), 849-859; discussion 863-844. PMID: 7944874.
- O'Connor, J.A., Wiffen, B., Diforti, M., Ferraro, L., Joseph, C., Kolliakou, A., Bonaccorso, S., Murray, R.M., David, A.S., 2013. Neuropsychological, clinical and cognitive insight predictors of outcome in a first episode psychosis study. *Schizophr Res* **149** (1-3), 70-76. PMID: 23815972.
- O'Donovan, M.C., Craddock, N., Norton, N., Williams, H., Peirce, T., Moskvina, V., Nikolov, I., Hamshere, M., Carroll, L., Georgieva, L., Dwyer, S., Holmans, P., Marchini, J.L., Spencer, C.C., Howie, B., Leung, H.T., Hartmann, A.M., Moller, H.J., Morris, D.W., Shi, Y., Feng, G., Hoffmann, P., Propping, P., Vasilescu, C., Maier, W., Rietschel, M., Zammit, S., Schumacher, J., Quinn, E.M., Schulze, T.G., Williams, N.M., Giegling, I., Iwata, N., Ikeda, M., Darvasi, A., Shifman, S., He, L., Duan, J., Sanders, A.R., Levinson, D.F., Gejman, P.V., Cichon, S., Nothen, M.M., Gill, M., Corvin, A., Rujescu, D., Kirov, G., Owen, M.J., Buccola, N.G., Mowry, B.J., Freedman, R., Amin, F., Black, D.W., Silverman, J.M., Byerley, W.F., Cloninger, C.R., Molecular Genetics of Schizophrenia Collaboration, 2008. Identification of loci associated with schizophrenia by genome-wide association and follow-up. *Nat Genet* **40** (9), 1053-1055. PMID: 18677311.
- Owens, S.F., Picchioni, M.M., Ettinger, U., McDonald, C., Walshe, M., Schmechtig, A., Murray, R.M., Rijdsdijk, F., Touloupoulou, T., 2012. Prefrontal deviations in function but not volume are putative endophenotypes for schizophrenia. *Brain* **135** (Pt 7), 2231-2244. PMID: 22693145.
- Owens, S.F., Picchioni, M.M., Rijdsdijk, F.V., Stahl, D., Vassos, E., Rodger, A.K., Collier, D.A., Murray, R.M., Touloupoulou, T., 2011a. Genetic overlap between episodic memory deficits and schizophrenia: Results from the Maudsley Twin Study. *Psychol Med* **41** (3), 521-532. PMID: 20459888.
- Owens, S.F., Rijdsdijk, F., Picchioni, M.M., Stahl, D., Nenadic, I., Murray, R.M., Touloupoulou, T., 2011b. Genetic overlap between schizophrenia and selective components of executive function. *Schizophr Res* **127** (1-3), 181-187. PMID: 21056927.
- Pasternak, O., Westin, C.F., Bouix, S., Seidman, L.J., Goldstein, J.M., Woo, T.U., Petryshen, T.L., Mesholam-Gately, R.I., McCarley, R.W., Kikinis, R., Shenton, M.E., Kubicki, M., 2012. Excessive extracellular volume reveals a neurodegenerative pattern in schizophrenia onset. *J Neurosci* **32** (48), 17365-17372. PMID: 23197727.
- PGC Schizophrenia Working Group, 2014. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511** (7510), 421-427. PMID: 25056061.
- Psychosis Endophenotypes International Consortium, Wellcome Trust Case-Control Consortium, Bramon, E., Pirinen, M., Strange, A., Lin, K., Freeman, C., Bellenguez, C., Su, Z., Band, G., Pearson, R., Vukcevic, D., Langford, C., Deloukas, P., Hunt, S., Gray, E., Dronov, S., Potter, S.C., Tashakkori-Ghanbaria, A., Edkins, S., Bumpstead, S.J., Arranz, M.J., Bakker, S., Bender, S., Bruggeman, R., Cahn, W., Chandler, D., Collier, D.A., Crespo-Facorro, B., Dazzan, P., de Haan, L., Di Forti, M., Dragovic, M., Giegling, I., Hall, J., Iyegbe, C., Jablensky, A., Kahn, R.S., Kalaydjieva, L., Kravariti, E., Lawrie, S., Linszen, D.H., Mata, I., McDonald, C., McIntosh, A., Myin-Germeys, I., Ophoff, R.A., Pariente, C.M., Paunio, T., Picchioni, M., Psychiatric Genomics Consortium, Ripke, S., Rujescu, D., Sauer, H., Shaikh, M., Sussmann, J., Suvisaari, J., Tosato, S., Touloupoulou, T., Van Os, J., Walshe, M., Weisbrod, M., Whalley, H., Wiersma, D., Blackwell, J.M., Brown, M.A., Casas, J.P., Corvin, A., Duncanson, A., Jankowski, J.A., Markus, H.S., Mathew, C.G., Palmer, C.N., Plomin, R., Rautanen, A., Sawcer, S.J., Trembath, R.C., Wood, N.W., Barroso, I., Peltonen, L., Lewis, C.M., Murray, R.M., Donnelly, P., Powell, J., Spencer, C.C., 2014. A genome-wide association analysis of a broad psychosis phenotype identifies three loci for further investigation. *Biol Psychiatry* **75** (5), 386-397. PMID: 23871474.
- Quan, M., Lee, S.H., Kubicki, M., Kikinis, Z., Rathi, Y., Seidman, L.J., Mesholam-Gately, R.I., Goldstein, J.M., McCarley, R.W., Shenton, M.E., Levitt, J.J., 2013. White matter tract abnormalities between rostral middle frontal gyrus, inferior frontal gyrus and striatum in first-episode schizophrenia. *Schizophr Res* **145** (1-3), 1-10. PMID: 23415471.
- Rais, M., Cahn, W., Schnack, H.G., Hulshoff Pol, H.E., Kahn, R.S., van Haren, N.E., 2012. Brain volume reductions in medication-naïve patients with schizophrenia in relation to intelligence quotient. *Psychol Med* **42** (9), 1847-1856. PMID: 22357376.
- Randolph, C., Tierney, M.C., Mohr, E., Chase, T.N., 1998. The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): Preliminary clinical validity. *J Clin Exp Neuropsychol* **20** (3), 310-319. PMID: 9845158.
- Reis Marques, T., Taylor, H., Chaddock, C., Dell'acqua, F., Handley, R., Reinders, A.A., Mondelli, V., Bonaccorso, S., Diforti, M., Simmons, A., David, A.S., Murray, R.M., Pariente, C.M., Kapur, S., Dazzan, P., 2014. White matter integrity as a predictor of response to treatment in first episode psychosis. *Brain* **137** (Pt 1), 172-182. PMID: 24253201.
- Reitan, R., 1958. Validity of the trail making test as indicator of organic brain damage. *Percept Mot Skills* **8** (271-276). PMID: 13263471.

- Reitan, R., 1985. Halstead-Reitan Neuropsychological Test Battery: Theory and Clinical Interpretation. Reitan Neuropsychology, Tucson, Arizona.
- Rey, A., 1964. L'examen clinique en psychologie. Presses Universitaires de France, Paris.
- Rice, J.P., Reich, T., Bucholz, K.K., Neuman, R.J., Fishman, R., Rochberg, N., Hesselbrock, V.M., Nurnberger, J.I., Jr., Schuckit, M.A., Begleiter, H., 1995. Comparison of direct interview and family history diagnoses of alcohol dependence. *Alcohol Clin Exp Res* **19** (4), 1018-1023. PMID: 7485811.
- Robertson, I.H., Manly, T., Andrade, J., Baddeley, B.T., Yiend, J., 1997. 'Oops!': Performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia* **35** (6), 747-758. PMID: 9204482.
- Roffman, J.L., Lambert, J.S., Achtyes, E., Macklin, E.A., Galendez, G.C., Raeke, L.H., Silverstein, N.J., Smoller, J.W., Hill, M., Goff, D.C., 2013. Randomized multicenter investigation of folate plus vitamin B12 supplementation in schizophrenia. *JAMA Psychiatry* **70** (5), 481-489. PMID: 23467813.
- Roffman, J.L., Nitenson, A.Z., Agam, Y., Isom, M., Friedman, J.S., Dyckman, K.A., Brohawn, D.G., Smoller, J.W., Goff, D.C., Manoach, D.S., 2011. A hypomethylating variant of MTHFR, 677C>T, blunts the neural response to errors in patients with schizophrenia and healthy individuals. *PLoS One* **6** (9), e25253. PMID: 21980405.
- Roffman, J.L., Weiss, A.P., Deckersbach, T., Freudenreich, O., Henderson, D.C., Purcell, S., Wong, D.H., Halsted, C.H., Goff, D.C., 2007. Effects of the methylenetetrahydrofolate reductase (MTHFR) C677T polymorphism on executive function in schizophrenia. *Schizophr Res* **92** (1-3), 181-188. PMID: 17344026.
- Roffman, J.L., Weiss, A.P., Deckersbach, T., Freudenreich, O., Henderson, D.C., Wong, D.H., Halsted, C.H., Goff, D.C., 2008a. Interactive effects of COMT Val108/158Met and MTHFR C677T on executive function in schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* **147B** (6), 990-995. PMID: 18186041.
- Roffman, J.L., Weiss, A.P., Purcell, S., Caffalett, C.A., Freudenreich, O., Henderson, D.C., Bottiglieri, T., Wong, D.H., Halsted, C.H., Goff, D.C., 2008b. Contribution of methylenetetrahydrofolate reductase (MTHFR) polymorphisms to negative symptoms in schizophrenia. *Biol Psychiatry* **63** (1), 42-48. PMID: 17543893.
- Ruff, R.M., Parker, S.B., 1993. Gender- and age-specific changes in motor speed and eye-hand coordination in adults: Normative values for the Finger Tapping and Grooved Pegboard Tests. *Percept Mot Skills* **76** (3 Pt 2), 1219-1230. PMID: 8337069.
- Russell, A.J., Munro, J., Jones, P.B., Hayward, P., Hemsley, D.R., Murray, R.M., 2000. The National Adult Reading Test as a measure of premorbid IQ in schizophrenia. *Br J Clin Psychol* **39** (Pt 3), 297-305. PMID: 11033751.
- Schmand, B., Bakker, D., Saan, R., Louman, J., 1991. De Nederlandse Leestest voor Volwassenen: Een maat voor het premorbide intelligentieniveau [The Dutch Reading Test for Adults: a measure of premorbid intelligence level]. *Tijdschr Gerontol Geriatr* **22** (1), 15-19. PMID: 1877068.
- Seidman, L.J., Breiter, H.C., Goodman, J.M., Goldstein, J.M., Woodruff, P.W., O'Craven, K., Savoy, R., Tsuang, M.T., Rosen, B.R., 1998. A functional magnetic resonance imaging study of auditory vigilance with low and high information processing demands. *Neuropsychology* **12** (4), 505-518. PMID: 9805320.
- Seidman, L.J., Cherkertzian, S., Goldstein, J.M., Agnew-Blais, J., Tsuang, M.T., Buka, S.L., 2013. Neuropsychological performance and family history in children at age 7 who develop adult schizophrenia or bipolar psychosis in the New England Family Studies. *Psychol Med* **43** (1), 119-131. PMID: 22575089.
- Seidman, L.J., Kremen, W.S., Koren, D., Faraone, S.V., Goldstein, J.M., Tsuang, M.T., 2002. A comparative profile analysis of neuropsychological functioning in patients with schizophrenia and bipolar psychoses. *Schizophr Res* **53** (1-2), 31-44. PMID: 11728836.
- Seidman, L.J., Thermenos, H.W., Poldrack, R.A., Peace, N.K., Koch, J.K., Faraone, S.V., Tsuang, M.T., 2006. Altered brain activation in dorsolateral prefrontal cortex in adolescents and young adults at genetic risk for schizophrenia: An fMRI study of working memory. *Schizophr Res* **85** (1-3), 58-72. PMID: 16632333.
- Seitz, J., Zuo, J.X., Lyall, A.E., Makris, N., Kikinis, Z., Bouix, S., Pasternak, O., Fredman, E., Dushin, J., Goldstein, J.M., Petryshen, T.L., Meshulam-Gately, R.I., Wojcik, J., McCarley, R.W., Seidman, L.J., Shenton, M.E., Koerte, I.K., Kubicki, M., 2016. Tractography analysis of 5 white matter bundles and their clinical and cognitive correlates in early-course schizophrenia. *Schizophr Bull* **42** (3), 762-771. PMID: 27009248.
- Shallice, T., 1982. Specific impairments of planning. *Philos Trans R Soc Lond B Biol Sci* **298** (1089), 199-209. PMID: 6125971.
- Sharma, T., Bilder, R., 2004. Standardisation and cross validation study of Cogtest - An automated neurocognitive battery for use in clinical trials. *Eur Neuropsychopharmacol* **14** (Supplement 3), S386. PMID: N/A.
- Simpson, G.M., Angus, J.W., 1970. A rating scale for extrapyramidal side effects. [Clinical Trial Comparative Study Controlled Clinical Trial]. *Acta Psychiatr Scand Suppl* **212**, 11-19. PMID: 4917967.
- Sivan, A.B., 1992. Benton Visual Retention Test, 5th ed. The Psychological Corporation, San Antonio, TX.
- Spitzer, R.L., Endicott, J., 1978. Schedule for Affective Disorders and Schizophrenia (SADS)-Lifetime version. New York State Psychiatric Institute, New York, NY.
- Stefansson, H., Ophoff, R.A., Steinberg, S., Andreassen, O.A., Cichon, S., Rujescu, D., Werge, T., Pietilainen, O.P., Mors, O., Mortensen, P.B., Sigurdsson, E., Gustafsson, O., Nyegaard, M., Tuulio-Henriksson, A., Ingason, A., Hansen, T., Suvisaari, J., Lonnqvist, J., Paunio, T., Borglum, A.D., Hartmann, A., Fink-Jensen, A., Nordentoft, M., Hougaard, D., Norgaard-Pedersen, B., Bottcher, Y., Olesen, J., Breuer, R., Moller, H.J., Giegling, I., Rasmussen, H.B., Timm, S., Mattheisen, M., Bitter, I., Rethelyi, J.M., Magnusdottir, B.B., Sigmundsson, T., Olason, P., Masson,

- G., Gulcher, J.R., Haraldsson, M., Fossdal, R., Thorgeirsson, T.E., Thorsteinsdottir, U., Ruggeri, M., Tosato, S., Franke, B., Strengman, E., Kiemene, L.A., Genetic Risk and Outcome in Psychosis (GROUPE), Melle, I., Djurovic, S., Abramova, L., Kaleda, V., Sanjuan, J., de Frutos, R., Bramon, E., Vassos, E., Fraser, G., Ettinger, U., Picchioni, M., Walker, N., Touloupoulou, T., Need, A.C., Ge, D., Yoon, J.L., Shianna, K.V., Freimer, N.B., Cantor, R.M., Murray, R., Kong, A., Golimbet, V., Carracedo, A., Arango, C., Costas, J., Jonsson, E.G., Terenius, L., Agartz, I., Petursson, H., Nothen, M.M., Rietschel, M., Matthews, P.M., Muglia, P., Peltonen, L., St Clair, D., Goldstein, D.B., Stefansson, K., Collier, D.A., 2009. Common variants conferring risk of schizophrenia. *Nature* **460** (7256), 744-747. PMID: 19571808.
- Stern, R.A., White, T., 2003. NAB administration, scoring, and interpretation manual. Psychological Assessment Resources, Inc., Lutz, FL.
- Stinissen, J., Willems, P.J., Coetsier, P., Hulsman, W.L.L., 1970. Manual for the Dutch Translated and Adapted Version of The Wechsler Adult Intelligence Scale (WAIS). Swets & Zeitlinger, Lisse, The Netherlands.
- Stroop, J.R., 1935. Studies of interference in serial verbal reactions. *J Exp Psychol* **18** (6), 643-662. PMID: N/A.
- Stroup, T.S., McEvoy, J.P., Swartz, M.S., Byerly, M.J., Glick, I.D., Canive, J.M., McGee, M.F., Simpson, G.M., Stevens, M.C., Lieberman, J.A., 2003. The National Institute of Mental Health Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) project: Schizophrenia trial design and protocol development. *Schizophr Bull* **29** (1), 15-31. PMID: 12908658.
- Sturm, W., Willmes, K., 1999. Verbaler Lerntest (VLT) [Verbal Learning Test]. Hogrefe, Göttingen, Germany.
- Sullivan, P.F., Lin, D., Tzeng, J.Y., van den Oord, E., Perkins, D., Stroup, T.S., Wagner, M., Lee, S., Wright, F.A., Zou, F., Liu, W., Downing, A.M., Lieberman, J., Close, S.L., 2008. Genomewide association for schizophrenia in the CATIE study: Results of stage 1. *Mol Psychiatry* **13** (6), 570-584. PMID: 18347602.
- Swagerman, S.C., de Geus, E.J., Kan, K.J., van Bergen, E., Nieuwboer, H.A., Koenis, M.M., Hulshoff Pol, H.E., Gur, R.E., Gur, R.C., Boomsma, D.I., 2016. The Computerized Neurocognitive Battery: Validation, aging effects, and heritability across cognitive domains. *Neuropsychology* **30** (1), 53-64. PMID: 26710096.
- Terwisscha van Scheltinga, A.F., Bakker, S.C., van Haren, N.E., Buizer-Voskamp, J.E., Boos, H.B., Vorstman, J., Cahn, W., Hulshoff Pol, H., Ophoff, R.A., Kahn, R.S., 2012. Association study of copy number variants with brain volume in schizophrenia patients and healthy controls. *Psychiatry Res* **200** (2-3), 1011-1013. PMID: 22551941.
- Terwisscha van Scheltinga, A.F., Bakker, S.C., van Haren, N.E., Derks, E.M., Buizer-Voskamp, J.E., Boos, H.B., Cahn, W., Hulshoff Pol, H.E., Ripke, S., Ophoff, R.A., Kahn, R.S., Psychiatric Genome-wide Association Study Consortium, 2013a. Genetic schizophrenia risk variants jointly modulate total brain and white matter volume. *Biol Psychiatry* **73** (6), 525-531. PMID: 23039932.
- Terwisscha van Scheltinga, A.F., Bakker, S.C., van Haren, N.E., Derks, E.M., Buizer-Voskamp, J.E., Cahn, W., Ripke, S., Psychiatric Genome-Wide Association Study Consortium, Ophoff, R.A., Kahn, R.S., 2013b. Schizophrenia genetic variants are not associated with intelligence. *Psychol Med* **43** (12), 2563-2570. PMID: 23410598.
- Tewes, U., 1991. Hamburg-Wechsler Intelligenztest für Erwachsene-Revision [WAIS-R, German version]. Hogrefe, Göttingen, Germany.
- The Cognition Group, 2006. Cogtest. Cogtest, Inc., Newark, DE.
- Theleritis, C., Fisher, H.L., Shafer, I., Winters, L., Stahl, D., Morgan, C., Dazzan, P., Breedvelt, J., Sambath, I., Vitoratou, S., Russo, M., Reichenberg, A., Falcone, M.A., Mondelli, V., O'Connor, J., David, A., McGuire, P., Pariante, C., Di Forti, M., Murray, R.M., Bonaccorso, S., 2014. Brain derived Neurotrophic Factor (BDNF) is associated with childhood abuse but not cognitive domains in first episode psychosis. *Schizophr Res* **159** (1), 56-61. PMID: 25171858.
- Torgesen, J.K., Wagner, R., Rashotte, C., 1999. Test of Word Reading Efficiency (TOWRE). ProEd Publishing Co., Austin, TX.
- Touloupoulou, T., Grech, A., Morris, R.G., Schulze, K., McDonald, C., Chapple, B., Rabe-Hesketh, S., Murray, R.M., 2004. The relationship between volumetric brain changes and cognitive function: A family study on schizophrenia. *Biol Psychiatry* **56** (6), 447-453. PMID: 15364043.
- Touloupoulou, T., Morris, R.G., Rabe-Hesketh, S., Murray, R.M., 2003. Selectivity of verbal memory deficit in schizophrenic patients and their relatives. *Am J Med Genet B Neuropsychiatr Genet* **116B** (1), 1-7. PMID: 12497605.
- Touloupoulou, T., Picchioni, M., Rijdsdijk, F., Hua-Hall, M., Ettinger, U., Sham, P., Murray, R., 2007. Substantial genetic overlap between neurocognition and schizophrenia: Genetic modeling in twin samples. *Arch Gen Psychiatry* **64** (12), 1348-1355. PMID: 18056542.
- Trampush, J.W., Yang, M.L., Yu, J., Knowles, E., Davies, G., Liewald, D.C., Starr, J.M., Djurovic, S., Melle, I., Sundet, K., Christoforou, A., Reinvang, I., DeRosse, P., Lundervold, A.J., Steen, V.M., Espeseth, T., Raikkonen, K., Widen, E., Palotie, A., Eriksson, J.G., Giegling, I., Konte, B., Roussos, P., Giakoumaki, S., Burdick, K.E., Payton, A., Ollier, W., Horan, M., Chiba-Falek, O., Attix, D.K., Need, A.C., Cirulli, E.T., Voineskos, A.N., Stefanis, N.C., Avramopoulos, D., Hatzimanolis, A., Arking, D.E., Smyrnis, N., Bilder, R.M., Freimer, N.A., Cannon, T.D., London, E., Poldrack, R.A., Sabb, F.W., Congdon, E., Conley, E.D., Scult, M.A., Dickinson, D., Straub, R.E., Donohoe, G., Morris, D., Corvin, A., Gill, M., Hariri, A.R., Weinberger, D.R., Pendleton, N., Bitsios, P., Rujescu, D., Lahti, J., Le Hellard, S., Keller, M.C., Andreassen, O.A., Deary, I.J., Glahn, D.C., Malhotra, A.K., Lencz, T., 2017. GWAS meta-

- analysis reveals novel loci and genetic correlates for general cognitive function: A report from the COGENT consortium. *Mol Psychiatry* **22** (3), 336-345. PMID: 28093568.
- van 't Wout, M., Aleman, A., Kessels, R.P., Laroi, F., Kahn, R.S., 2004. Emotional processing in a non-clinical psychosis-prone sample. *Schizophr Res* **68** (2-3), 271-281. PMID: 15099609.
- van Erp, T.G., Hibar, D.P., Rasmussen, J.M., Glahn, D.C., Pearlson, G.D., Andreassen, O.A., Agartz, I., Westlye, L.T., Haukvik, U.K., Dale, A.M., Melle, I., Hartberg, C.B., Gruber, O., Kraemer, B., Zilles, D., Donohoe, G., Kelly, S., McDonald, C., Morris, D.W., Cannon, D.M., Corvin, A., Machielsen, M.W., Koenders, L., de Haan, L., Veltman, D.J., Satterthwaite, T.D., Wolf, D.H., Gur, R.C., Gur, R.E., Potkin, S.G., Mathalon, D.H., Mueller, B.A., Preda, A., Macciardi, F., Ehrlich, S., Walton, E., Hass, J., Calhoun, V.D., Bockholt, H.J., Sponheim, S.R., Shoemaker, J.M., van Haren, N.E., Pol, H.E., Ophoff, R.A., Kahn, R.S., Roiz-Santanez, R., Crespo-Facorro, B., Wang, L., Alpert, K.I., Jonsson, E.G., Dimitrova, R., Bois, C., Whalley, H.C., McIntosh, A.M., Lawrie, S.M., Hashimoto, R., Thompson, P.M., Turner, J.A., 2016. Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry* **21** (4), 547-553. PMID: 26283641.
- Vassos, E., Di Forti, M., Coleman, J., Iyegbe, C., Prata, D., Euesden, J., O'Reilly, P., Curtis, C., Kolliakou, A., Patel, H., Newhouse, S., Traylor, M., Ajnakina, O., Mondelli, V., Marques, T.R., Gardner-Sood, P., Aitchison, K.J., Powell, J., Atakan, Z., Greenwood, K.E., Smith, S., Ismail, K., Pariante, C., Gaughran, F., Dazzan, P., Markus, H.S., David, A.S., Lewis, C.M., Murray, R.M., Breen, G., 2017. An examination of polygenic score risk prediction in individuals with first-episode psychosis. *Biol Psychiatry* **81** (6), 470-477. PMID: 27765268.
- Versmissen, D., Janssen, I., Myin-Germeys, I., Mengelers, R., Campo, J.A., van Os, J., Krabbendam, L., 2008. Evidence for a relationship between mentalising deficits and paranoia over the psychosis continuum. *Schizophr Res* **99** (1-3), 103-110. PMID: 17936589.
- Vollema, M.G., Ormel, J., 2000. The reliability of the Structured Interview for Schizotypy-Revised. *Schizophr Bull* **26** (3), 619-629. PMID: 10993402.
- Wagner, R.K., Torgesen, J.K., Rashotte, C.A., 1999. Comprehensive test of phonological processing. PRO-ED inc., Austin, TX.
- Walters, J.T., Corvin, A., Owen, M.J., Williams, H., Dragovic, M., Quinn, E.M., Judge, R., Smith, D.J., Norton, N., Giegling, I., Hartmann, A.M., Moller, H.J., Muglia, P., Moskvina, V., Dwyer, S., O'Donoghue, T., Morar, B., Cooper, M., Chandler, D., Jablensky, A., Gill, M., Kaladjeva, L., Morris, D.W., O'Donovan, M.C., Rujescu, D., Donohoe, G., 2010. Psychosis susceptibility gene ZNF804A and cognitive performance in schizophrenia. *Arch Gen Psychiatry* **67** (7), 692-700. PMID: 20603450.
- Walters, J.T., Rujescu, D., Franke, B., Giegling, I., Vasquez, A.A., Hargreaves, A., Russo, G., Morris, D.W., Hoogman, M., Da Costa, A., Moskvina, V., Fernandez, G., Gill, M., Corvin, A., O'Donovan, M.C., Donohoe, G., Owen, M.J., 2013. The role of the major histocompatibility complex region in cognition and brain structure: A schizophrenia GWAS follow-up. *Am J Psychiatry* **170** (8), 877-885. PMID: 23903335.
- Wechsler, D., 1974. Wechsler Adult Intelligence Scale-Revised (WAIS-R). The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1981. Examiner's Manual: Wechsler Adult Intelligence Scale-Revised (WAIS-R). The Psychological Corporation, New York, NY.
- Wechsler, D., 1987. Wechsler Memory Scale-Revised (WMS-R) Manual. The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1991. Wechsler Intelligence Scale for Children-Third Edition (WISC-III). The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1997a. Wechsler Adult Intelligence Scale-Third Edition (WAIS-III). The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1997b. Wechsler Memory Scale-Third Edition (WMS-III). The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 1999. Wechsler Abbreviated Scale of Intelligence (WASI). The Psychological Corporation, San Antonio, TX.
- Wechsler, D., 2000. Wechsler Adult Intelligence Scale – Third Edition. Nederlandstalige Bewerking. Technische Handleiding. Pearson Assessment and information B.V., Amsterdam, Netherlands.
- Wechsler, D., 2001. Wechsler Test of Adult Reading. Harcourt Assessment, San Antonio, TX.
- Wechsler, D., 2005. Wechsler Adult Intelligence Scale-Third Edition, UK Edition (WAIS-III-UK). Psychological Corporation Ltd, London, UK.
- Wheeler, A.L., Chakravarty, M.M., Lerch, J.P., Pipitone, J., Daskalakis, Z.J., Rajji, T.K., Mulsant, B.H., Voineskos, A.N., 2014. Disrupted prefrontal interhemispheric structural coupling in schizophrenia related to working memory performance. *Schizophr Bull* **40** (4), 914-924. PMID: 23873858.
- Wilkinson, G.S., 1993. Wide Range Achievement Test-Third Edition (WRAT-3). Jastak Association, Wilmington, DE.
- Wilkinson, G.S., Robertson, G.R., 2006a. Manual for the Wide Range Achievement Test-Fourth Edition (WRAT-4). Psychological Assessment Resources, Lutz, FL.
- Wilkinson, G.S., Robertson, G.R., 2006b. Wide Range Achievement Test 4 (WRAT-4) Professional Manual. Psychological Assessment Resources, Lutz, FL.

Wing, J.K., Babor, T., Brugha, T., Burke, J., Cooper, J.E., Giel, R., Jablenski, A., Regier, D., Sartorius, N., 1990. SCAN. Schedules for Clinical Assessment in Neuropsychiatry. *Arch Gen Psychiatry* **47** (6), 589-593. PMID: 2190539.

Woodcock, R.W., McGrew, K.S., Mather, N., 2001. Woodcock-Johnson III Test. Riverside Publishing Company, Itasca, IL.

World Health Organization, 1992. The ICD-10 classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines. World Health Organisation, Geneva, Switzerland.

Supplementary Table 1. Description of the GENUS Consortium Sample Collection (subset with ge

| Acronym | Sample | Site |
|-------------------|--|----------------------|
| CAMH | Centre for Addiction and Mental Health | Toronto, Canada |
| CATIE | Clinical Antipsychotic Trials of Intervention Effectiveness | Multi-site, USA |
| CIDAR/VA | Longwood Study of Psychosis consortium / VA Healthcare System | Boston, USA |
| COGS-UK | Cognition and Genetics in Schizophrenia & Bipolar Disorder | Cardiff, UK |
| GAP | Genetics and Psychosis First-Episode Study | London, UK |
| IMH-SIGNRP | Institute of Mental Health - Singapore Imaging Genetics and Neuropsychological Research in Psychosis | Singapore |
| IMH-STCRP | Institute of Mental Health - Singapore Translational and Clinical Research in Psychosis | Singapore |
| KCL-MFS | King's College London - Maudsley Family Study | London, UK |
| KCL-MTS | King's College London - Maudsley Twin Study | London, UK |
| L&R | Language and Risk in Schizophrenia | Boston, USA |
| MCIC | Mind Clinical Imaging Consortium | Multi-site, USA |
| MGH | Massachusetts General Hospital | Boston, USA |
| NEFS | New England Family Study | Boston, USA |
| PAGES | Phenomics and Genomics Sample | Munich, Germany |
| PHRS | Pittsburgh High Risk Study | Pittsburgh, USA |
| TCD/NUIG | Trinity College Dublin/ National University of Ireland, Galway | Multi-site, Ireland |
| UMCU-SZ1 | University Medical Center Utrecht - Schizophrenia Study 1 | Utrecht, Netherlands |
| UMCU-SZ2 | University Medical Center Utrecht - Schizophrenia Study 2 | Utrecht, Netherlands |
| ZHH | Zucker Hillside Hospital | New York, USA |
| Totals | | |

Data in this table are based on genotyped participants only.

All samples that acquired T1 MRI scans also acquired DTI scans except the PHRS and UMCU-SZ1

Population ancestry determined from genetic data (where available) or self report.

Eur = European-derived ancestry; FHR = familial high-risk.

notype data).

| GWAS Array | Neuropsychological data | | | | |
|--|--------------------------------|-----------------|------------|-------------|------------|
| | Patients | Controls | FHR | Male | Eur |
| | (N) | (N) | (N) | (%) | (%) |
| Illumina OmniExpress | 90 | 107 | 0 | 54.8 | 76.1 |
| Affymetrix 500K; Perlegen's custom 164K chip | 741 | 0 | 0 | 73.6 | 54.7 |
| Illumina OmniExpress | 74 | 106 | 5 | 68.1 | 60.0 |
| Illumina Infinium OmniExpressExome-8 | 529 | 0 | 0 | 60.7 | 97.3 |
| Illumina HumanCore-24 Exome BeadChip | 136 | 127 | 0 | 59.3 | 46.8 |
| Illumina Human OmniZhongHua-8; Illumina Human1M-Duo; Affymetrix 6.0 | 104 | 3 | 0 | 51.4 | 0 |
| Illumina HumanOmniZhongHua-8 BeadChip | 419 | 956 | 0 | 53.0 | 0 |
| Affymetrix 6.0 | 10 | 12 | 19 | 56.1 | 95.1 |
| Affymetrix 6.0 | 18 | 65 | 5 | 35.2 | 100 |
| Illumina Infinium PsychArray | 0 | 30 | 41 | 36.6 | 74.7 |
| Illumina HumanOmni1-Quad BeadChip | 80 | 66 | 0 | 69.9 | 75.3 |
| Illumina Infinium PsychArray | 423 | 0 | 0 | 72.1 | 68.8 |
| Illumina Infinium PsychArray | 75 | 125 | 24 | 47.3 | 86.2 |
| Illumina OmniExpress; Illumina HumanHap300 | 210 | 1,341 | 0 | 50.0 | 99.6 |
| Illumina Infinium PsychArray | 0 | 18 | 43 | 47.5 | 41.0 |
| Affymetrix 6.0; Illumina HumanCore Exome | 826 | 165 | 0 | 63.3 | 99.9 |
| Illumina HumanHap550; Illumina Infinium OmniExpressExome-8 | 50 | 96 | 0 | 68.5 | 98.6 |
| Illumina HumanHap550; Affymetrix 6.0; Illumina Infinium OmniExpressExome-8 | 189 | 115 | 168 | 58.7 | 97.5 |
| Illumina OmniExpress | 0 | 219 | 0 | 49.3 | 100 |
| | 3,974 | 3,551 | 305 | 56.6 | 72.2 |

y.
 scans except the PHRS and UMCU-SZ1
 (available) or self report.

| <i>T1-weighted structural MRI data</i> | | | | |
|--|-----------------|------------|-------------|------------|
| <i>Patients</i> | <i>Controls</i> | <i>FHR</i> | <i>Male</i> | <i>Eur</i> |
| <i>(N)</i> | <i>(N)</i> | <i>(N)</i> | <i>(%)</i> | <i>(%)</i> |
| 89 | 115 | 0 | 55.4 | 76.5 |
| --- | --- | --- | --- | --- |
| 66 | 100 | 5 | 67.3 | 60.2 |
| --- | --- | --- | --- | --- |
| 67 | 35 | 0 | 60.8 | 37.3 |
| 159 | 3 | 0 | 59.9 | 0 |
| --- | --- | --- | --- | --- |
| --- | --- | --- | --- | --- |
| 7 | 20 | 3 | 50.0 | 100 |
| 0 | 31 | 44 | 36.0 | 76.0 |
| 82 | 68 | 0 | 69.3 | 74.7 |
| 44 | 110 | 0 | 62.3 | 72.7 |
| 66 | 126 | 14 | 46.6 | 84.0 |
| --- | --- | --- | --- | --- |
| 0 | 11 | 33 | 45.5 | 45.5 |
| 134 | 189 | 0 | 54.5 | 99.7 |
| 74 | 101 | 0 | 69.7 | 98.3 |
| 154 | 103 | 151 | 58.8 | 97.6 |
| --- | --- | --- | --- | --- |
| 942 | 1,012 | 250 | 56.6 | 71.0 |

Supplementary Table 2. Clinical and demographic characteristics of the GENUS Consortium Sam

| | Patients | |
|--|-----------------|---|
| | N | Mean \pm SD (Range) |
| Age (years) | 4,101 | 39.8 \pm 12.1 (13-82) |
| Education Level (years) | 3,740 | 12.2 \pm 2.5 (1-22) |
| Premorbid IQ | 2,644 | 96.2 \pm 15.7 (44-145) |
| Current IQ | 1,478 | 93.9 \pm 18.4 (47-155) |
| Illness Duration (years) | 3,296 | 15.0 \pm 11.5 (<1-58) |
| Age at Onset (years) | 3,278 | 24.5 \pm 8.3 (1-62) |
| Global Assessment of Functioning | 1,396 | 60.4 \pm 15.7 (20-100) |
| PANSS Positive symptoms | 2,507 | 16.7 \pm 7.3 (7-47) |
| PANSS Negative Symptoms | 2,504 | 17.1 \pm 7.2 (7-42) |
| PANSS General Symptoms | 2,510 | 32.7 \pm 12.0 (0-93) |
| SAPS Positive Symptoms | 1,149 | 9.1 \pm 13.3 (0-121) |
| SANS Negative Symptoms | 887 | 24.4 \pm 20.3 (0-103) |
| Antipsychotic dose – current CPZEQ | 2,727 | 579.9 \pm 414.8 (0-5,000) |
| Antipsychotic dose – lifetime average CPZEQ | 1,355 | 534.6 \pm 360.8 (0-2,125) |
| | N | % |
| Sex (male / female; % male) | 2,721 / 1,381 | 66.3 |
| Antipsychotic medication | | |
| Atypical | 1,691 | 48.2 |
| Typical | 350 | 10.0 |
| Both Typical and Atypical | 517 | 14.7 |
| Naïve / None | 375 | 10.7 |
| Unknown Class | 236 | 6.7 |
| No information | 338 | 9.6 |
| Diagnosis | | |
| Schizophrenia | 3,321 | 81.0 |
| Schizoaffective Disorder | 340 | 8.3 |
| Schizophreniform Disorder | 67 | 1.6 |
| Bipolar Psychosis | 207 | 5.1 |
| Other Psychosis | 150 | 3.7 |
| Psychosis Unknown Type | 17 | 0.4 |
| Ethnicity | | |
| European | 2,835 | 69.1 |
| East Asian | 624 | 15.2 |
| African | 454 | 11.1 |
| American (Predominantly Latino) | 138 | 3.4 |
| South Asian | 42 | 1 |
| Mixed | 11 | 0.3 |
| Handedness (right / other; % right-handed) | 1,508 / 151 | 90.1 |

Data in this table are based on genotyped participants only. CPZEQ = chlorpromazine 100 mg equivalent; df = deg

ple Collection (subset with genotype data).

[illegible]

p

<1x10⁻¹⁰

<1x10⁻¹⁰

<1x10⁻¹⁰

<1x10⁻¹⁰

<1x10⁻¹⁰

<1x10⁻¹⁰

0.38

Supplementary Table 3. Sample size by neuropsychological test across GENUS Consortium samp

| Domain | Attention / Processing Speed | | | | | | |
|------------|------------------------------|--------------------------|---------------------|--------|---------------------------------------|----------------------------------|--------------------------|
| | Digit Symbol Coding | | | TMT-A | Verbal Fluency | | |
| | BACS Symbol Coding* | WAIS Digit Symbol-Coding | RBANS Symbol Coding | TMT-A* | Semantic/ Category Fluency - Animals* | Semantic/ Category Fluency - Any | Phonemic/ Letter Fluency |
| CAMH | | | x | x | RBANS | | COWAT |
| CATIE | | | | | x | x | COWAT |
| CIDAR/VA | x | | | x | MCCB | x | |
| COGS-UK | x | | | x | MCCB | x | |
| GAP | | III | | x | x | x | COWAT |
| IMH-SIGNRP | x | | | | | BACS | COWAT |
| IMH-STCRP | x | | | | BACS | BACS | |
| KCL-MFS | | R | | x | | | |
| KCL-MTS | | III-UK | | x | x | x | COWAT |
| L&R | x | | | x | MCCB | x | |
| MCIC | | | | x | D-KEFS | D-KEFS | D-KEFS |
| MGH | | III | | | x | x | COWAT |
| NEFS | | R | | | | | COWAT |
| PAGES | | R-DE | | x | x | x | x |
| PHRS | | | | | MAE | MAE | MAE |
| TCD/ NUIG | | | | x | | | COWAT |
| UMCU-SZ1 | | | | | MAE | MAE | MAE |
| UMCU-SZ2 | | III-NL | | | | | |
| ZHH | x | | | x | MCCB | x | COWAT |
| N Patients | 1,470 | 1,909 | 116 | 1,549 | 3,139 | 3,414 | 2,626 |
| N Controls | 1,405 | 2,156 | 114 | 1,116 | 2,445 | 2,611 | 1,647 |
| N FHR | 50 | 346 | 0 | 196 | 247 | 247 | 229 |

| | | | | | | | |
|---------|-------|-------|-----|-------|-------|-------|-------|
| N Total | 2,925 | 4,411 | 230 | 2,861 | 5,831 | 6,272 | 4,502 |
|---------|-------|-------|-----|-------|-------|-------|-------|

*MATRICS test.

References for all neuropsychological tests are provided in the Supplemental Materials.

Abbreviations: 128-P, 128-C = 128-card paper, computerized version; 64-P, 64-C = 64-card paper, computerized version; Cognition in Schizophrenia; BVMT-R = Brief Visuospatial Memory Test-Revised; BVRT = Benton Visual Retention Test; CMS = Children's Memory Scale; CNB = Computerized Neurocognitive Battery; CPT(-IP) = Continuous Performance Test (-Intercept) version; D-KEFS = Delis-Kaplan Executive Function System; FHR = Familial High Risk; FigMem = Figural Memory Test; L-O = Test of Line Orientation; NAB = Neuropsychological Assessment Battery; NL = Dutch version; PAL = Paired Associates Test; NPS = Neuropsychological Status; RST = Response Shifting Task; RVIP = Rapid Visual Information Processing; SART = Sustained Attention to Response Test; Cambridge; SWM = Spatial Working Memory; TOL = Tower of London; TMT-A, B = Trail Making Test Part A, B; UK = United Kingdom; PAL = Paired Associates; VisRep = Visual Reproduction; VLMT = Verbal Learning and Memory Test; VLT = Verbal Learning Test; WISC-III = Wechsler Intelligence Scale for Children; WMS = Wechsler Memory Scale; WAIS = Wechsler Adult Intelligence Scale.

les (genotyped + ungenotyped).

| Attention / Vigilance | | | | | | | |
|------------------------|------|----------------|---------|---------|---------|---------|---------|
| Continuous Performance | | | | | | | |
| CPT-IP* | ACPT | CANTAB RVIP | 3-7 CPT | A-X CPT | 1-9 CPT | H-Q CPT | X-Y CPT |
| | | | | | | | |
| X | | | | | | | |
| X | X | | | | | | |
| X | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| X | | | | | | | |
| | | | X | | | | |
| | | | X | | | | |
| X | X | | | | | | |
| | | | | | | | |
| X | | | | | | | |
| | X | | | | | | |
| | | | X | | | | |
| X | | | | X | | | |
| X | | | | | X | | X |
| | | | | | | X | |
| | | | | | | X | |
| X | | | | | | | |
| 2,337 | 122 | 44 | 201 | 0 | 83 | 246 | 85 |
| 1,410 | 256 | 74 | 411 | 22 | 60 | 198 | 42 |
| 119 | 83 | 21 | 0 | 66 | 0 | 211 | 0 |

| | | | | | | | |
|-------|-----|-----|-----|----|-----|-----|-----|
| 3,866 | 461 | 139 | 612 | 88 | 143 | 655 | 127 |
|-------|-----|-----|-----|----|-----|-----|-----|

rd paper, computerized version; ACPT = Auditory CPT; AVLTL = Auditory Verbal Learning Test; BACS = Brief Assessment of Benton Visual Retention Test; CANTAB = Cambridge Neuropsychological Test Automated Battery; CFT = Complex Figure Test; CPT = Continuous Performance Test (Identical Pairs); CVLT(-SF) = California Verbal Learning Test (Short Form); DE = German; FigMem = Figural Memory; HVLTL = Hopkins Verbal Learning Test; IDED = Intra-Extra Dimensional Set Shifting; JOLO = Judgment of Line Orientation; PAL = Paired Associates Learning; RAN = Rapid Automatized Naming; RBANS = Repeatable Battery for the Assessment of Neuropsychological Status; SART = Sustained Attention to Response Task; SDRT = Spatial Delayed Response Task; SOC = Stockings of Color; Making Test Part A, B; UK = British version; UMD = University of Maryland; VisMemSpan = Visual Memory Span; VisPA = Visual Pattern Association Test; VLT = Verbal Learning Test; VOLT = Visual Object Learning Test; WASI = Wechsler Abbreviated Scale of Intelligence; WAIS = Wechsler Adult Intelligence Scale; WCST = Wisconsin Card Sorting Test

| Working Memory – verbal | | | | | | Working Memory – non-verbal | |
|-------------------------|-------------------------------|------------------|---------------------|-----------------------|---------------------|-----------------------------|-----------------------------|
| Letter-Number Span | | Other | | | | Spatial Span | |
| UMD Letter-Number Span* | WAIS Letter-Number Sequencing | RBANS Digit Span | WAIS/WMS Digit Span | BACS Digit Sequencing | WAIS/WMS Arithmetic | WMS-III Spatial Span* | WMS-R-UK Visual Memory Span |
| x | | x | | | | | |
| x | | | | | | | |
| x | | | | | | x | |
| x | | | | | | x | |
| | | | III | | | x | |
| | | | | x | | | |
| | | | | x | | | |
| | | | R | | R | | |
| | III-UK | | III-UK | | III-UK | | x |
| x | | | | | | x | |
| | III | | | | | | |
| | III | | III | | III | | |
| | | | R | | | | |
| | | | R-DE | | R-DE | | |
| | | | | | | | |
| | III | | | | | | |
| | | | | | | | |
| | | | | | III-NL | | |
| x | | | R | | | x | |
| 1,734 | 1,168 | 565 | 115 | 637 | 1,001 | 1,024 | 73 |
| 468 | 616 | 1,075 | 141 | 1,889 | 1,752 | 429 | 181 |
| 49 | 30 | 0 | 0 | 114 | 314 | 50 | 26 |

| | | | | | | | |
|-------|-------|-------|-----|-------|-------|-------|-----|
| 2,251 | 1,814 | 1,640 | 256 | 2,640 | 3,067 | 1,503 | 280 |
|-------|-------|-------|-----|-------|-------|-------|-----|

BACS = Brief Assessment of
ery; CFT = Complex Figure Test;
(Short Form); DE = German
nal Set Shifting; JOLO = Judgment
Battery for the Assessment of
onse Task; SOC = Stockings of
ual Memory Span; VisPA = Visual
riated Scale of Intelligence; WISC =

| Working Memory – non-verbal | | | | Verbal Learning | | | |
|-----------------------------|------------|--------|-------------|--------------------|--------|---------|-------|
| Other | | | | Word List Learning | | | |
| SDRT | CANTAB SWM | n-back | Cogtest SWM | HVLT* | CVLT | CVLT-SF | BACS |
| | | | | | | | |
| | x | | | x | | | |
| | | | | R | | | |
| | | | | R | | | |
| | | x | | | | | |
| | | | | | | | x |
| | | | | | | | x |
| | | x | | | | | |
| | | x | | | | | |
| | | | | R | | | |
| | | | | R | | | |
| | | | | | | x | |
| | | | | | x / II | | |
| | | | x | | | | |
| | | | x | | | | |
| | | x | x | | | | x |
| | | | | | | | I-NL |
| | | | | | | | |
| | | | x | | R | | |
| 656 | 792 | 268 | 0 | 1,737 | 562 | 100 | 558 |
| 0 | 448 | 433 | 48 | 426 | 227 | 36 | 1,074 |
| 0 | 21 | 0 | 68 | 50 | 32 | 0 | 0 |

| | | | | | | | |
|-----|-------|-----|-----|-------|-----|-----|-------|
| 656 | 1,261 | 701 | 116 | 2,213 | 821 | 136 | 1,632 |
|-----|-------|-----|-----|-------|-----|-----|-------|

| Verbal Learning & Memory | | | | | | | | |
|--------------------------|------|-----------------|------|--|--------------------------|--------------------------|------------------|---|
| Word List Learning | | | | | Story Recall | | | Other |
| RBANS | AVLT | Cogtest WLMT | VLMT | | WMS Logical Memory | CMS Logical Memory | RBANS Stories | WMS Verbal Paired Associate s |
| x | | | | | | | x | |
| | | | | | | | | |
| | | | | | III | x | | |
| | | | | | | | | |
| | | | | | III | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | R | | | R |
| | | | | | R-UK | | | R-UK |
| | | | | | III | | | |
| | | | | | III | | | |
| | | | | | | | | |
| | | | | | R/ III | | | |
| | | | x | | R-DE | | | R-DE |
| | | x | | | | | | |
| | | | | | III | | | |
| | | | | | | | | |
| | x | | | | | | | |
| | | | | | | | | |
| 116 | 213 | 0 | 206 | | 1,330 | 6 | 116 | 388 |
| 141 | 142 | 49 | 426 | | 863 | 13 | 141 | 705 |
| 0 | 231 | 71 | 0 | | 82 | 0 | 0 | 177 |

| | | | | | | | |
|-----|-----|-----|-----|-------|----|-----|-------|
| 257 | 586 | 120 | 632 | 2,275 | 19 | 257 | 1,270 |
|-----|-----|-----|-----|-------|----|-----|-------|

| Visual Learning & Memory | | | | | | | |
|--------------------------|---------------------------|--------------------------|--------------------------------|--------------|------|---------|---|
| BVMT-R* | RBANS Figure Recall | WMS Figural Memory | WMS Visual Reproductio n | WMS Faces | BVRT | Rey CFT | WMS Visual Paired Associate s |
| | | x | | | | | |
| | x | | | | | | |
| | x | | | | | | |
| | | | III | | | | |
| | | | | | | | |
| | | | R | | | | |
| | | | R-UK | | | | R-UK |
| | x | | | | | | |
| | | | R | III | x | | |
| | | | | | | | |
| | | | | III | | x | |
| | | R-DE | R-DE | | | | R-DE |
| | | | | | | | |
| | | | | III | | | |
| | | | R-NL | | | | |
| | | | | | | | |
| | x | | | | | | |
| 897 | 115 | 206 | 655 | 712 | 112 | 32 | 279 |
| 328 | 140 | 427 | 950 | 458 | 95 | 116 | 608 |
| 48 | 0 | 0 | 228 | 25 | 0 | 31 | 26 |

| | | | | | | | |
|-------|-----|-----|-------|-------|-----|-----|-----|
| 1,273 | 255 | 633 | 1,833 | 1,195 | 207 | 179 | 913 |
|-------|-----|-----|-------|-------|-----|-----|-----|

Reasoning / Problem

| | | Planning | | | | | |
|---------------|----------|---------------|-------------------|----------|-----|--------|---------------|
| CANTAB PAL | CNB VOLT | NAB Mazes* | WISC-III Mazes | BACS TOL | TOL | TOL-DE | CANTAB SOC |
| | | | | | | | |
| | | | X | | | | |
| | | X | | | | | |
| | | X | | | | | |
| | | | | | | | X |
| | | | | | X | | |
| | | | | | X | | |
| | | | | | | | |
| | | | | | | | |
| | | X | | | | | |
| | | | | | | X | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | X | |
| | X | | | | | | |
| X | | | | | | | |
| | | | | | | | |
| | | X | | | | | |
| | | X | | | | | |
| 608 | 0 | 903 | 715 | 551 | 93 | 202 | 109 |
| 242 | 50 | 328 | 0 | 1,075 | 84 | 427 | 117 |
| 0 | 56 | 50 | 0 | 0 | 0 | 0 | 0 |

| | | | | | | | |
|-----|-----|-------|-----|-------|-----|-----|-----|
| 850 | 106 | 1,281 | 715 | 1,626 | 177 | 629 | 226 |
|-----|-----|-------|-----|-------|-----|-----|-----|

Reasoning / Problem Solving

| | Set Shifting + Planning | Cognitive Flexibility / Set Shifting | | | Response Inhibition | | | |
|--|-------------------------|--------------------------------------|---------------|-------------|---------------------|-----|------|------------------|
| | TMT-B | WCST 64-Card | WCST 128-Card | CANTAB IDED | Stroop | RST | SART | Cogtest Go-No-Go |
| | x | | | | x | | | |
| | | 64-C | | | | | | |
| | | 64-C | | | | | | |
| | | | | | | | | |
| | x | | | | | | | |
| | | | | | | | | |
| | | 64-P | | | | | | |
| | x | | | x | | | | |
| | x | | | x | | | | |
| | | 64-C | | | | | | |
| | x | | | | | | | |
| | | 64-C | 128-C | | | | | |
| | | | 128-P | | x | | | |
| | x | | 128-C | | | | | |
| | | | 128-P | | | | | x |
| | x | | | x | | | x | |
| | | | | | x | | | |
| | | | | | | x | | |
| | x | 128-P | | | x | | | |
| | 836 | 1,001 | 375 | 522 | 201 | 194 | 513 | 0 |
| | 1,408 | 132 | 703 | 285 | 404 | 129 | 182 | 49 |
| | 145 | 57 | 77 | 21 | 0 | 209 | 0 | 70 |

| | | | | | | | |
|-------|-------|-------|-----|-----|-----|-----|-----|
| 2,389 | 1,190 | 1,155 | 828 | 605 | 532 | 695 | 119 |
|-------|-------|-------|-----|-----|-----|-----|-----|

Visuo-spatial Ability

| WAIS Block Design | WAIS Matrix Reasoning | WAIS Object Assembly | WAIS Picture Arrangement | WAIS Picture Completion | RBANS JOLO | Benton JOLO |
|-------------------|-----------------------|----------------------|--------------------------|-------------------------|------------|-------------|
| | | | | | x | |
| WASI | | | | | | |
| III | III | | | | | |
| | | | | | | |
| | WASI | | | | | x |
| R | | R | R | R | | |
| III-UK | | III-UK | III-UK | III-UK | | |
| WASI | | | | | | |
| III | | | | | | |
| III | III | III | III | III | | |
| R | | | R | | | |
| R-DE | | R-DE | R-DE | R-DE | | |
| | | | | | | |
| III-R-UK | III-R-UK | | | | | |
| III-R-NL | | | III-R-NL | | | |
| III-NL | | | | | | |
| | | | | | | |
| 2,260 | 1,149 | 311 | 456 | 330 | 116 | 417 |
| 2,744 | 1,385 | 1,340 | 1,607 | 1,347 | 141 | 1,012 |
| 522 | 0 | 0 | 33 | 0 | 0 | 0 |

| | | | | | | |
|-------|-------|-------|-------|-------|-----|-------|
| 5,526 | 2,534 | 1,651 | 2,096 | 1,677 | 257 | 1,429 |
|-------|-------|-------|-------|-------|-----|-------|

| | | Verbal Ability | | | | |
|----------------------|-----------------|--------------------|-----------------------|---------------------|----------------------|--------------------|
| | | | | | | |
| RBANS Figure Copy | Rey CFT Copy | WAIS Vocabulary | WAIS Comprehension | WAIS Information | WAIS Similarities | D-KEFS Proverbs |
| x | | | | | | |
| | | WASI | | | | x |
| | | | | III | | |
| | | | | | | |
| | | R | R | R | R | |
| | | III-UK | III-UK | III-UK | III-UK | |
| | | WASI | | | | x |
| | | III | | | III | |
| | | III | III | III | III | |
| | x | R | R | R | | |
| | | R-DE | R-DE | R-DE | R-DE | |
| | | | | | | |
| | | III-R-UK | | | III-R-UK | |
| | | III-R-NL | III-R-NL | | | |
| | | | | III-NL | | |
| | | | | | | |
| 116 | 32 | 1,754 | 456 | 1,027 | 1,140 | 66 |
| 140 | 116 | 2,425 | 1,607 | 1,645 | 1,935 | 128 |
| 0 | 33 | 285 | 33 | 233 | 253 | 48 |

| | | | | | | |
|-----|-----|-------|-------|-------|-------|-----|
| 256 | 181 | 4,464 | 2,096 | 2,905 | 3,328 | 242 |
|-----|-----|-------|-------|-------|-------|-----|

